

U.S. FISH AND WILDLIFE SERVICE
DIVISION OF ENVIRONMENTAL QUALITY
REGION 6

**ENVIRONMENTAL CONTAMINANTS ASSOCIATED WITH A
SWINE CONCENTRATED ANIMAL FEEDING OPERATION
AND IMPLICATIONS FOR McMURTREY NATIONAL
WILDLIFE REFUGE.**



U.S. Fish and Wildlife Service
Nebraska Field Office
203 West Second Street
Grand Island, Nebraska 68801
July, 2004

**ENVIRONMENTAL CONTAMINANTS ASSOCIATED WITH A SWINE
CONCENTRATED ANIMAL FEEDING OPERATIONS AND IMPLICATIONS FOR
MCMURTREY NATIONAL WILDLIFE REFUGE.**

Prepared by
Matthew S. Schwarz¹
Kathy R. Echols²
Mark J. Wolcott³
Karen J. Nelson⁴

¹U.S. Fish and Wildlife Service
Division of Environmental Quality
Nebraska Field Office
Grand Island, Nebraska 68801

²U.S. Geological Survey
Biological Resources Division
Columbia Environmental Research Center
Columbia, Missouri 65210

³U.S. Geological Survey
Biological Resources Division
National Wildlife Health Center
Madison, Wisconsin 53711

⁴U.S. Fish and Wildlife Service
Division of Environmental Quality
Montana Field Office
Helena, Montana 59601

U.S. Fish and Wildlife Service
Division of Environmental Quality
Region 6
DEC ID:6N45
FFS:200060006

ACKNOWLEDGEMENTS

This research would not have been possible without the support and partnership of Hastings Pork. Owen Nelson, the Hastings Pork Farm Manager, was particularly interested in whether the created wetlands posed any problems for migratory waterfowl and was instrumental to this research by providing insight into farm operations. The authors also thank those that contributed time and effort in completing this study. Steve Moran contributed funding for water quality analyses. Dr. Dan Snow conducted the analyses of antibiotics in water and sediment samples. Brenda Berlowski and Heather Gutzman analyzed all samples for bacterial pathogens and assisted with sample collections. Jeff Runge monitored water quality during the summer and fall of 2000. Christina Lydick, Jeff Runge, Steve Lydick, Jeanette O'Neal, and John Meisner assisted with several aspects of the field work. Beth Goldowitz assisted with the logistics behind invertebrate diversity sampling and analysis. Jeff Drahota performed the waterfowl brood surveys. Dr. David Marx provided technical support on statistical analyses and performed the proc mixed analysis in SAS®. Dr. James Rosowski provided technical support to identify algae. Andy Bishop helped construct site map figures. John Cochnar, Christina Lydick, Brenda Berlowski, Dan Martin, Craig Giggelman, and Dr. Dan Snow reviewed report drafts. Larry Gamble reviewed the report draft and coordinated project funding and sample submission.

TABLE OF CONTENTS

| | |
|---------------------------------------|-----|
| ACKNOWLEDGEMENTS | iii |
| ABSTRACT | vi |
| LIST OF TABLES | vii |
| LIST OF FIGURES | ix |
| INTRODUCTION | 1 |
| Nature and Scope of the Problem | 1 |
| Site Description and History | 4 |
| Research Justification and Objectives | 7 |
| Contaminants Associated with CAFOs | 7 |
| Trace Elements | 8 |
| Salts | 12 |
| Nutrients | 13 |
| Algal Toxins | 14 |
| Bacterial Pathogens | 14 |
| Antibiotics | 16 |
| Hormones | 18 |
| METHODS | 20 |
| Algal Toxins | 20 |
| Antibiotics | 21 |
| Bacterial Pathogens | 21 |
| Hormones | 24 |
| Waterfowl Use | 24 |
| Invertebrate Diversity | 24 |
| Water Quality | 25 |
| Trace Elements | 25 |

| | |
|------------------------------------|----|
| Statistical Analyses | 27 |
| RESULTS | 29 |
| Algal Toxins | 29 |
| Antibiotics | 30 |
| Bacterial Pathogens | 33 |
| Hormones | 36 |
| Waterfowl Use | 38 |
| Invertebrate Analysis | 38 |
| Water Quality | 40 |
| Trace Elements | 43 |
| Sediment | 43 |
| Water | 43 |
| Invertebrates | 44 |
| DISCUSSION | 51 |
| Trace Elements | 52 |
| Antibiotics | 54 |
| Bacterial Pathogens | 55 |
| Hormones | 57 |
| Water Quality | 58 |
| Uncertainty Analysis and Data Gaps | 59 |
| Ongoing Research | 59 |
| Future Research Needs | 60 |
| Recommendations | 61 |
| Conclusions | 62 |
| REFERENCES | 64 |
| APPENDIX: Additional Tables | 75 |

ABSTRACT

Waste generated by concentrated animal feeding operations (CAFOs) may contain a variety of contaminants including nutrients, pathogens, trace elements, antibiotics, and hormones. In 2000, the U.S. Fish and Wildlife Service began to characterize CAFO contaminants in lagoons, canals, and created wetlands operated by Hastings Pork, a large swine CAFO adjacent to McMurtrey National Wildlife Refuge (McMurtrey) in Clay County, Nebraska. The created wetlands were designed to attract waterfowl; therefore, the primary purpose of this research was to evaluate whether migratory waterfowl were likely exposed to CAFO contaminants. A secondary research objective was to determine if created wetland water was suitable as a supplementary water source for McMurtrey. Wetlands created from swine wastewater effluent had 5-50 fold greater concentrations of phosphorus, ammonia, and total nitrogen and 2-3 fold greater salinity compared to control sites. Cyanobacteria (*Microcystis* spp.) were abundant in the created wetlands and microcystin toxins were detected in concentrated water samples. Tetracycline, macrolide, and diterpene antibiotics were detected in lagoon and canal sediment and water samples; however, in the created wetlands only oxytetracycline was detected (once in sediment at 41 nanograms per gram). Concentrations of 17- β estradiol and testosterone in CAFO wastewater (n=4) exceeded toxicity thresholds for aquatic life. Fecal coliform and streptococci counts in water (n=38) generally exhibited a decreasing gradient with lagoons > canals > created wetlands > McMurtrey. Bacteria (*Salmonella* spp. and *Yersinia enterocolitica*) were recovered in the created wetlands but not McMurtrey. Created wetland invertebrate communities were dominated by chironomid species and had lower taxa diversity when compared to McMurtrey. Eutrophication of created wetlands may represent the greatest health threat to waterfowl by creating an environment conducive to cyanobacteria blooms and outbreaks of avian botulism and avian cholera. Trace elements from swine waste will likely continue to accumulate in the created wetlands over time, leading to an increased risk of exposure to wetland biota. Research is ongoing and includes use of sentinel mallards (*Anas platyrhynchos*) to further evaluate the need to decrease concentrations of CAFO contaminants in swine wastewater before it is used to create waterfowl habitat.

LIST OF TABLES

| | | |
|-----|---|----|
| 1 | Summary of samples collected for trace element analyses at Hastings Pork and McMurtrey Marsh, 2000 to 2001. | 28 |
| 2 | Total microcystin (MC) concentrations in algal blooms from the Hastings Pork created wetlands, Clay County, Nebraska, 2000. | 29 |
| 3 | Mean annual fecal coliforms and fecal streptococci colony forming units per 100 ml of water from sites at Hastings Pork and McMurtrey Marsh, Clay County, Nebraska, 2000. | 33 |
| 4 | Concentrations of 17- β estradiol and testosterone in water samples collected from McMurtrey Marsh and Hastings Pork created wetlands, canals and lagoons in Clay County, Nebraska, 2000, compared to those published by other studies. | 37 |
| 5 | Total number of individuals by invertebrate order found in the spring, summer and fall of 2000 from sites at McMurtrey Marsh and created wetlands 1 and 5. | 39 |
| 6 | Results from Proc Mixed in the Statistical Analysis System (SAS) for trace elements in sediments from Rainwater Basin wetlands and Hastings Pork lagoons, canals, and created wetlands in Clay County, Nebraska | 45 |
| 7 | Mean (\pm SE) trace element concentrations in sediments from the Hastings Pork created wetlands and Rainwater Basin wetlands compared to U.S. western background concentrations and effects thresholds. | 47 |
| 8 | Total recoverable concentrations (mg/L) of trace elements in water samples collected from McMurtrey Marsh and Hastings Pork canals and created wetlands, Clay County, Nebraska. | 48 |
| 9 | Results from Proc Mixed in the Statistical Analysis System (SAS) for trace elements in chironomids from Hastings Pork, Nebraska; Stillwater National Wildlife Refuge (NWR), Nevada; and the Sun River Irrigation Project Area, Montana. | 50 |
| A.1 | Summary statistics for concentrations of trace elements in water samples from Rainwater Basin wetlands and Hastings Pork created wetlands and canals, Clay County, Nebraska. | 76 |

| | | |
|------|--|----|
| A.2 | Monthly counts of fecal coliforms and fecal streptococci colony forming units per 100 ml of water from sites at Hastings Pork and McMurtrey Marsh, Clay County, Nebraska, 2000. | 77 |
| A.3 | <i>Salmonella</i> species (serotype) recovered from Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000. | 78 |
| A.4 | <i>Yersinia</i> species recovered from McMurtrey Marsh and Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000. | 79 |
| A.5. | Concentrations (mg/L) of trace elements in water samples collected from Waterfowl Production Areas (federally managed wetlands) in the Rainwater Basin, Clay County, Nebraska, 1992-2001. | 80 |
| A.6. | Concentrations (mg/L) of trace elements in sediment samples collected from Waterfowl Production Areas (federally managed wetlands) in the Rainwater Basin, Clay County, Nebraska, 1992-2001. | 81 |
| A.7. | Antibiotic concentrations in sediment (ng/g) and water (µg/g) from Hastings Pork created wetlands, canals and lagoons, Clay County, Nebraska, 2000. | 82 |
| A.8 | Concentrations (µg/g) of trace elements in invertebrate samples collected from Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000 - 2001. | 83 |
| A.9 | Concentrations (µg/g) of trace elements in sediment samples collected from McMurtrey Marsh and Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000. | 84 |
| A.10 | Concentrations (mg/L) of trace elements in water samples collected from McMurtrey Marsh and Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000. | 87 |

LIST OF FIGURES

| | | |
|----|--|----|
| 1 | Location of the Rainwater Basin Region at the focal point of a spring migration corridor used by millions of ducks, geese and other migratory birds annually. | 3 |
| 2 | Location of the Nebraska Rainwater Basin and the study site in Clay County, Nebraska. | 5 |
| 3 | Location of the Hastings Pork created wetlands, McMurtrey National Wildlife Refuge (NWR), and McMurtrey Marsh. | 6 |
| 4 | Locations of samples collected for identification of bacteria pathogens at Hastings Pork and McMurtrey NWR, Clay County, Nebraska, 2000 | 23 |
| 5 | Mean (\pm S.E.) concentrations of antibiotics in sediments from lagoons and canals at Hastings Pork, Clay County, NE, 2000. | 31 |
| 6 | Mean (\pm S.E.) concentrations of antibiotics in water from lagoons and canals at Hastings Pork, Clay County, NE, 2000. | 32 |
| 7 | Mean (\pm S.E.) fecal coliform and fecal streptococci counts for water samples collected from created wetlands (CW), canals, lagoons, and McMurtrey Marsh (MM) in Clay County, Nebraska, 2000. | 34 |
| 8 | Number of isolates of <i>Yersinia</i> and <i>Salmonella</i> species from McMurtrey Marsh and Hastings Pork created wetlands, canals and lagoons, Clay County, NE, 2000. | 36 |
| 9 | Shannon-Weaver taxa diversity for benthic invertebrates collected in the spring, summer, and fall from McMurtrey Marsh, Created Wetland 1, and Created Wetland 5 in Clay County, NE, 2000. | 40 |
| 10 | Mean (\pm SE) measurements of pH, dissolved oxygen, and specific conductivity in the Hastings Pork created wetlands, McMurtrey Marsh and Rainwater Basin (RWB) wetlands in Clay County, NE, 2000. | 41 |

| | | |
|----|---|----|
| 11 | Mean (\pm SE) concentrations of ammonia, total Kjeldahl nitrogen, and total phosphorus in Hastings Park created wetlands, McMurtrey Marsh and Rainwater Basin wetlands in Clay County, NE, 2000. | 42 |
| 12 | Mean (\pm SE) concentrations of lead (Pb) in sediments from Rainwater Basin wetlands and Hayden Thompson created wetlands, canals, and lagoons, Clay County, NE, 2000. | 46 |
| 13 | Mean (\pm SE) concentrations of boron (B), magnesium (Mg) and manganese (Mn) in water from Rainwater Basin wetlands and Hastings Park created wetlands and canals, Clay County, Nebraska, 2000. | 49 |

INTRODUCTION

Nature and Scope of the Problem

The trend in livestock operations in the U.S., with fewer operations and increased numbers of animals per operation, has created a concern that the animal wastes from these facilities may represent an increased risk to the environment. According to the U.S. Environmental Protection Agency (USEPA), the number of animal feeding operations for hogs in the top ten production states in the U.S. decreased from over 300,000 to under 150,000 from 1978 to 1992; however, during this same period, the average number of hogs per operation increased by 134 percent (USEPA, 2001). A similar trend is occurring in Nebraska where, according to the Nebraska Agricultural Statistics Service (NASS), the average number of hogs per farm more than doubled from 368 to 853 in 1992 and 2001, respectively (NASS, 1996 and 2002). This concentration of animals and their waste is not without consequence to the environment. According to the Nebraska Department of Environmental Quality (NDEQ), the number of fish kills attributed to livestock wastes in Nebraska increased from 5 in 1998 and 1999 to 11 in 2002 and 2003 (NDEQ, unpublished data).

A “large” swine concentrated animal feeding operation (CAFO) is defined by USEPA as a facility with more than 2,500 swine, each weighing over 25 kilograms (kg), in confinement (USEPA, 2003). Pollutants associated with these facilities include trace elements, salts, nutrients, cyanobacteria toxins, bacterial pathogens, hormones, and antibiotics (USEPA, 2003). These pollutants can enter rivers, streams, and wetlands by spills, lagoon ruptures, field run-off, and contaminated groundwater. Adverse effects of swine lagoon breaches to fish and wildlife are well documented (Mallin et al., 1997; Burkholder et al., 1997; Williams, 1998; Denn, 1999). A 25 million gallon swine wastewater spill in 1995 from a CAFO in North Carolina killed 10 million fish and closed 364,000 acres of coastal wetlands to shellfishing (Williams, 1998). In Missouri, swine CAFOs were designated as the biggest culprit in polluting 150 miles of Missouri's streams, causing 61 fish kills and killing more than 500,000 fish (Denn, 1999). Land

application of animal manure also can lead to the accumulation of heavy metals and phosphorus in soil. When soil adsorption sites become limited, the ability to bind excess phosphorus and metals decreases and the soil changes from a sink to a source for the transport of these elements to surface run-off (Sharpley et al., 1999). Buffalo Lake National Wildlife Refuge in Texas experienced this in the 1960's and 1970's when large fish kills on the refuge were attributed to field run-off and discharges from cattle feedlots within the refuge's watershed (Baker et al., 1998). Water quality degradation eventually led to the draining of Buffalo Lake. Wetlands at the refuge now receive well water to compensate for the nutrient loading from run-off (Baker et al., 1998).

The Rainwater Basin (RWB), named for its abundance of natural wetlands, is increasingly at risk from CAFO run-off. The RWB region encompasses more than 4,200 square miles within 17 counties of south-central Nebraska and is recognized as the focal point of a spring migration corridor used by millions waterfowl and shorebirds annually (RWB JV, 2000; Figure 1). During the spring waterfowl migration, the RWB hosts approximately 90 percent of the mid-continent population of greater white-fronted geese (*Ansu albifrons*), 50 percent of the mid-continent population of mallards (*Anas platyrhynchos*), and 30 percent of the continent population of northern pintails (*Anas acuta*) (Benning 1987; Bortner et al., 1991). Threatened and endangered species including the whooping crane (*Grus americana*), bald eagle (*Haliaeetus leucocephalus*), and piping plover (*Charadrius melodus*) also have been observed at RWB wetlands (Wally Jobman, U.S. Fish and Wildlife Service Wildlife Biologist, pers. comm., 2004). The U.S. Fish and Wildlife Service (Service) Rainwater Basin Wetland Management District (RWB-WMD) manages 61 Waterfowl Production Areas (WPAs) that range in size from 38 to 1,989 acres (USFWS, 2001). Between 131 to 166 CAFOs are in operation within the RWB-WMD, and at least five WPAs have CAFOs in their watershed (Nebraska Department of Environmental Quality, unpublished data, 2002).

In 1991, the RWB was identified by the North American Waterfowl Management Plan as waterfowl habitat of major concern in North America and received Joint Venture status. The overall goal of the Rainwater Basin Joint Venture (RWB JV) is to restore and

maintain wetland habitat within the RWB. The RWBJV promotes sustainable agricultural practices to reduce the level of certain chemicals and other environmental contaminants from entering wetlands (Gersib et al., 1992).




Figure 1. Location of the Rainwater Basin Region at the focal point of a spring migration corridor used by millions of ducks, geese and other migratory birds annually. Note: figure taken from the Rainwater Basin Joint Venture Evaluation Plan (RWBJV, 2000).

Site Description and History

The study site was located in Clay County, Nebraska, within the RWB (Figure 2) and included Hastings Pork and McMurtrey National Wildlife Refuge (NWR). Hastings Pork is located on what was formerly a Naval Ammunition Depot. Since the 1960s, Hastings Pork has utilized the area for livestock and crop production. About 260 bunkers that were formerly used by the Navy to store munitions are now used for swine production. These bunkers house approximately 64,000 swine and around 1.5 million liters of water per day is used to flush them out. In addition, an estimated 325,000 liters of swine urine and manure slurry are generated daily (based on calculations for swine between 36-55 kg of body weight; Fraser 1991), for a total of 1.8 million liters of wastewater per day. In an effort to utilize this wastewater, a partnership between Hastings Pork and the RWBJV resulted in the creation of seven wetlands (known as the Hayden Thompson wetlands and referred to herein as the created wetlands) totaling 17 acres on Hastings Pork property. These wetlands receive swine wastewater effluent from lagoons by a canal system, with a distance of delivery ranging from less than a mile to 5 miles. The created wetlands were designed to provide waterfowl habitat and were not intended to treat swine-waste effluent; therefore, the Service and Hastings Pork formed a partnership to evaluate whether migratory birds attracted to the created wetlands may be exposed to contaminants and disease pathogens. The Service is also concerned that waterfowl may transmit disease pathogens from Hastings Pork to nearby habitats. McMurtrey Marsh is located within McMurtrey NWR and is located approximately 1 mile east of the created wetlands (Figure 3). McMurtrey NWR contains 650 acres of wetlands and 400 acres of upland habitat and is managed primarily for migratory waterfowl.





Figure 2. Location of the Nebraska Rainwater Basin and the study site in Clay County, Nebraska.



Scale: |-----|
1 mile

Figure 3. Location of the Hastings Pork created wetlands, McMurtrey National Wildlife Refuge (NWR), and McMurtrey Marsh. Note: See title page for a picture of the created wetlands and bunkers.

Research Justification and Objectives

To our knowledge, there have been no studies that have evaluated risk to waterfowl that utilize wetlands created from primary treated CAFO wastewater. However, many contaminants associated with CAFOs are known to be potentially toxic to waterfowl. For example, selenium or boron are directly toxic to waterfowl and can cause reproductive failure (USDI, 1998). Other contaminants in swine waste may not be very toxic to waterfowl directly (e.g., ammonia and nutrients), but have the potential to adversely affect waterfowl populations indirectly through habitat modification. Excessive nutrient loading can cause wetland eutrophication and may result in an environment with less foraging potential (Gaiser and Lang, 1998) or an environment more conducive to harmful cyanobacterial blooms or disease pathogens (Crowder and Bristow, 1988; USGS, 1999a).

The purpose of this research was to evaluate whether migratory waterfowl that utilize the created wetlands and McMurtrey NWR are likely exposed to CAFO contaminants including trace elements, nutrients, salts, cyanobacterial toxins, bacterial pathogens, hormones, and antibiotics. Sampling was performed to evaluate movement of contaminants from lagoons through canals and to the created wetlands. Contaminants were tested in water and sediments from a control site (McMurtrey Marsh), and in invertebrates, water, and sediments from the created wetlands, canals, and lagoons. In addition to the exposure assessment, taxa richness of invertebrate communities among McMurtrey Marsh and the created wetlands were compared to evaluate the effect of swine waste on the wetlands' function as waterfowl habitat.

Contaminants Associated with CAFOs

Trace elements, salts, nutrients, cyanobacterial toxins, bacterial pathogens, hormones, and antibiotics are swine CAFO pollutants of concern that may adversely affect natural ecosystems (USEPA, 2003). Background information on the toxicity of swine CAFO contaminants and their potential to adversely affect waterfowl and their habitat (e.g., plant and invertebrate food items) is presented below.

Trace elements

Trace elements in hog manure of environmental concern include aluminum, arsenic, boron, cadmium, copper, iron, lead, manganese, molybdenum, nickel, selenium, and zinc (USEPA, 2003). Many of these trace elements (e.g., selenium, copper, and zinc) are ingredients in hog feed. Copper (Cu) is added to swine feed to promote growth and control disease, and zinc (Zn) is commonly added to Cu-enriched rations to ameliorate Cu toxicity to swine (Payne et al., 1988). Trace elements from CAFOs may accumulate in sediments, water, and biota to concentrations that are toxic to plants and lead to reproductive impairment, poor body condition, and immune system dysfunction in animals (Stubbs and Cathey, 1999).

Aluminum (Al) toxicity and bioavailability to aquatic biota is largely dependent on its solubility and generally increases as pH decreases (Gensemer and Playle, 1999). Aluminum bioavailability and toxicity at a pH greater than 7.0 is largely unknown; however, at a pH below 5.5, Al can be toxic to many plant species (Sparling et al., 1997). Concentrations of 400 to 500 micrograms per liter ($\mu\text{g/L}$) of Al in water within a pH range of 4.0 to 4.3 had negligible effects on mortality in amphipods, snails, or insect larvae (Sparling et al., 1997). Waterfowl are most likely exposed to Al by dietary uptake and dietary concentrations less than ($<$) 1000 mg/kg are not considered harmful (Sparling et al., 1997). There is no evidence of aluminum bioaccumulation in aquatic invertebrates (Gensemer and Playle, 1999); however, there is a potential for food items including invertebrates, tadpoles, and a few species of plants, to have sufficient concentrations of Al to be toxic to avian species (Sparling et al., 1997).

Arsenic (As) toxicity and bioavailability depends on its chemical speciation, with trivalent (As^{+3}) generally being more toxic than pentavalent (As^{+5}). Pentavalent As predominates in most aquatic environments where it is relatively persistent and may bioaccumulate in aquatic organisms (USDI, 1998). Concentrations of As^{+5} in water ranging from 1 to 15.2 $\mu\text{g/L}$ have been reported to disrupt aquatic ecosystems by inhibiting the growth of certain aquatic plants (Sanders and Cibik, 1985 cited in Eisler, 1988a). Mallard ducklings fed As^{+5} at 30 mg/kg dry weight (dw) for 10 weeks exhibited

a reduced growth rate (Camardese et al., 1990). Adult mallards exhibited reduced body and liver mass, delayed onset of egg laying, decreased whole egg mass, and egg shell thinning when fed sodium arsenate at 400 mg/kg dw (Stanley et al., 1994).

Boron (B) toxicity is most often observed in plants where it functions as an essential trace element for growth and development (USDI, 1998). Boron in soil can cause toxic effects to plants at concentrations > 5 mg/L (Gupta et al., 1985 cited in USDI, 1998). Studies on the toxicity of B to aquatic invertebrates are lacking (USDI, 1998); however, adverse reproductive effects in the water flea (*Daphnia magna*) can occur at B concentrations of 13.6 mg/L (Gerisch, 1984). Waterborne concentrations of B greater than 450 mg/L can cause adverse reproductive effects in mallards and concentrations ranging from 30 mg/L to 300 mg/L can cause reduced weight gain in mallard ducklings (USDI, 1998).

Cadmium (Cd) is a known teratogen and mutagen, and can cause severe deleterious effects to fish and wildlife (Eisler, 1985). Freshwater biota is the most sensitive group. Concentrations of 0.8 to 9.9 mg/L Cd in water are lethal to several species of aquatic insects and crustaceans (Eisler, 1985). The kidney is the critical organ in avian species chronically exposed to cadmium, and young birds may be more susceptible to kidney damage than adults (Furness, 1996). Mallard ducklings given 20 mg/kg Cd for 90 days developed kidney damage (Cain et al., 1983) whereas adult mallards exhibited kidney damage at 200 mg/kg Cd (White et al., 1978).

Copper (Cu) toxicity mechanisms include free radical production, alteration in activities of several enzymes, and interference with metallothionein synthesis. Excess cellular copper leads to lipid peroxidation as superoxide radicals are created during the oxidation of Cu^{+1} to Cu^{+2} at the cellular membrane (Eisler, 1998a). Copper concentrations of 34 mg/kg in sediment rarely impair benthic invertebrate survival or reproduction, whereas concentrations above 270 mg/kg generally do (Long et al., 1995; Hull and Suter, 1994; USDI, 1998). Copper concentrations in soil greater than 25 to 50 mg/kg are toxic to sensitive plants (Demayo et al., 1982a). Copper is relatively nontoxic

to birds and mammals in comparison to aquatic invertebrates, plants, and fish as copper homeostasis in birds and mammals is well regulated by metallothionein (USDI, 1998).

Iron (Fe) can disrupt aquatic ecosystems by decreasing species diversity and abundance of periphyton and benthic invertebrates (Vuori, 1995; Dickman and Rygiel, 1998). Macroinvertebrates exposed to high levels of Fe, manganese, and concurrent blooms of iron-depositing bacteria (*Leptothrix ochracea*) exhibited varying responses including mortality by direct toxic effects and/or smothering, behavioral avoidance of bacterial-coated substrates, and an inability to feed on the bacteria (Wellnitz et al., 1994). Mallards ingesting eight BB size tungsten-iron shot over a 30-day period did not exhibit any adverse health effects (Kelly et al., 1998). No apparent adverse effects were observed in Mute Swans (*Cygnus olor*) after they received a diet containing moderately polluted sediments with high concentrations of Al, Fe, vanadium (V), and barium (Ba) for six weeks (Beyer et al., 2000).

Lead (Pb) targets kidney, bone, the central nervous system, and the hematopoietic system and can cause adverse biochemical, histopathological, neuropsychological, fetotoxic, teratogenic, immunotoxic, and reproductive effects (Eisler, 1988b; Goyer and Clarkson, 2001). *Daphnia magna* exhibit decreased reproduction in water with Pb concentrations as low as 1 µg/L (Vighi, 1981). Freshwater algae can bioconcentrate Pb to high levels (Eisler, 1988b). Algae exposed to Pb concentrations of 5 µg/L in water exhibited a bioconcentration factor of 92,000 (Demayo et al. 1982b). Ingestion of spent Pb shot has caused considerable mortality of migratory waterfowl and other birds, including raptors that eat waterfowl killed or wounded by hunters (Eisler, 1988b). Exposure to Pb in ways other than ingestion of Pb shot (or ingestible Pb objects such as sinkers) are unlikely to cause clinical signs of Pb poisoning in birds (Eisler, 1988b). Mallard hatchlings are apparently tolerant to Pb, as they do not exhibit adverse effects on growth at dietary levels of 500 mg Pb/kg, or survival at 2,000 mg Pb/kg (Eisler, 1988b).

Manganese (Mn) is considered the least toxic of the trace elements for poultry and mammals (Pais and Jones, 1997). However, the cycling of manganese between Mn⁺² and

Mn⁺³ may be potentially deleterious to biological systems because it can involve the generation of free radicals (Goyer and Clarkson, 2001).

Molybdenum (Mo) is relatively nontoxic to aquatic organisms and plants (USDI, 1998). Molybdenum toxicity is largely dependent on interactions with copper. A low copper-to-molybdenum ratio (<2), rather than the dietary Mo concentration, is the primary determinant of an organism's susceptibility to Mo poisoning (USDI, 1998). Aquatic organisms and plants generally do not exhibit adverse effects on growth or survival at water Mo concentrations < 50 mg/L; however, the same aquatic plants bioconcentrate Mo to levels potentially toxic to organisms that feed on them (USDI, 1998). Bioconcentration factors of 628; 3,300; and 3,570 have been reported for freshwater algae, cyanobacteria, and periphyton, respectively (USDI, 1998). Literature on Mo toxicity to wild birds is lacking; however, adverse effects from Mo exposure to domestic chickens include reduced growth at dietary concentrations of 200 to 300 mg/kg, decreased egg production at 500 mg/kg, and decreased survival at 6,000 mg/kg (Eisler, 1989).

Nickel (Ni) toxicity mechanisms include oxidative damage to DNA and proteins and the inhibition of cellular antioxidant defenses (Rodriguez et al., 1996 cited in Eisler, 1998b). Sensitive species of aquatic organisms are adversely affected at nominal waterborne concentrations of 11-113 mg Ni²⁺/L (Eisler, 1998b). Nickel compounds typically have a low hazard when administered orally (NAS, 1975 cited in Eisler 1998b; USEPA, 1980). Mallards fed diets containing 800 mg Ni/kg ration for 90 days exhibited metabolic upset and altered bone densities and mallards fed 1,200 mg Ni/kg exhibited reduced growth and survival (Cain and Pafford 1981; Eastin and O'Shea 1981).

Selenium (Se) is one of the most toxic trace elements with a narrow margin of safety between toxicity and dietary deficiency. Nutritionally optimal dietary Se exposure is generally reported as 0.1 to 0.3 mg/kg dw whereas thresholds for dietary toxicity in animals range from 2 to 5 mg/kg dw (USDI, 1998). The toxic effects of both Se deficiency and excess are similar and include reproductive depression, anemia, weight loss, and immune dysfunction (Koller and Exon, 1986; USDI, 1998). Vertebrates are

generally much more susceptible to Se toxicity than are most plants and invertebrates; therefore, the direct toxic effects of consuming Se-contaminated plants are believed to be more important than indirect ecological effects from changes in plant communities (USDI, 1998). Reproductive impairment (e.g. reduced hatchability and teratogenesis) can result in birds with diets containing as little as 3 to 8 mg Se per kg (Wilber, 1980; Heinz, 1996; USDI, 1998). Dabbling duck species are among the most Se sensitive waterbird species (USDI, 1998). The concentration of Se in duck eggs estimated to cause teratogenesis in duck eggs is 23 mg/kg dw (Skorupa 1998, USDI, 1998). The potential for Se to bioaccumulate and adversely affect sensitive species (including waterfowl) at waterborne concentrations less than 5 µg/L has resulted in the Service's request for the USEPA to develop a new chronic aquatic life water quality criterion of 2 µg/L.

Zinc is an essential element for all living organisms and is generally more toxic to aquatic invertebrates and plants than birds and mammals due to homeostatic control by metallothionein (USDI, 1998). Decreased growth rate in invertebrates has been reported for Zn concentrations > 10 µg/L and increased mortality at concentrations > 80 µg/L (Hatakeyama, 1989 cited in USDI, 1998; Eisler, 1993). Zinc poisoning in birds is indicated when liver concentrations exceed 2,100 mg/kg dw (Eisler, 1993). Mallards exposed to dietary Zn concentrations of 3,000 mg/kg dw for 30 days exhibited leg paralysis and decreased food consumption (Eisler, 1993).

Salts

Salinity is a measure of the mass of dissolved salts in a given mass of solution (USDI, 1998), and can be determined by measuring conductivity and then converting to parts per thousand (ppt). Salinity is acutely toxic to amphipods at a concentration of 22 ppt and to daphnia at 8 to 11 ppt (USDI, 1998). Waterfowl hatched in moderate to high saline environments and without access to fresh drinking water exhibit decreased growth, development, and survival rates (Stolley et al., 1999). Saline-induced mortality of ducklings or goslings generally happens before day 6 of life, after which the nasal salt glands are functional (Stolley et al., 1999). Salinity > 20 ppt is uniformly fatal to 48-

hour-old mallard and black duck ducklings (Barnes and Nudds, 1991). Salt glands collected from fully grown mallard and black ducks increased in size with increasing age and salinity, and hypertrophied to a maximum size at 1 percent NaCl, indicating a failure to regulate salts at a salinity > 10 ppt (Barnes and Nudds, 1991).

Nutrients

The primary nutrients released from hog manure are nitrogen, phosphorus, and potassium (USEPA, 2003). Nitrogen and phosphorus are high in livestock manure and can cause eutrophication in aquatic ecosystems. The average feeder hog will excrete 11 kg of nitrogen and 6 kg of phosphorus in one year (Fraser, 1991). Documented adverse effects to aquatic ecosystems that may develop following eutrophication include increased biomass of phytoplankton, shifts in the phytoplankton community to bloom-forming species that are toxic or inedible, decreased invertebrate and plant species diversity, and oxygen depletion (USEPA, 2003). Increased algal growth can disrupt aquatic ecosystems by consuming dissolved carbon dioxide and increasing pH (USEPA, 2003). Amphibian declines have been attributed, in part, to nitrite (NO_2^-) toxicity. Adverse effects of nitrate (NO_3^-) and nitrite exposure to five species of amphibian larvae included reduced feeding activity, less vigorous swimming, disequilibrium, paralysis, abnormalities, edemas, and death (Marco et al., 1999). These adverse effects increase with dose and, although sensitivity is different among species, all species showed 15-day LC50s (i.e., the concentration in water that is lethal to 50 percent of the test species in 15 days) lower than 2 mg nitrite per liter (Marco et al., 1999). Cascades frog tadpoles (*Rana cascadae*) exposed to 3.5 mg nitrite per liter exhibited retarded development and emerged at an earlier developmental stage (Marco and Blaustein, 1999). There are no known cases of acute toxicity to waterfowl from exposure to aqueous nitrogen or phosphorus; however, a die-off of 250 herring gulls (*Larus argentatus*) and ring-billed gulls (*Larus delawarensis*) in 1984 was attributed to ingestion of fertilizer waste containing 1,730 parts per million (ppm) nitrite (Ley, 1986).

Algal Toxins

Toxins produced by cyanobacteria (blue-green algae) have caused mortality among a variety of wildlife populations including amphibians, fish, snakes, waterfowl, raptors, deer, muskrats, fox, squirrels, skunks, mink, bats, and bees (Carmichael, 1992). A common class of toxins produced by cyanobacteria is the hepatotoxic microcystins. There are 52 microcystin variants, all of which share a similar acute toxic mechanism (Carmichael, 1997). Microcystins inhibit the protein phosphatases needed to control liver blood circulation, resulting in extensive hemorrhaging in the liver (Sivonen, 1996). Microcystin-LR (MC-LR) is one of the most common microcystin variants and rodent toxicity tests indicate that it is also one of the most toxic (Carmichael, 1997). The acute lethal dose to 50 percent of the treated population (LD50) for mice given an intraperitoneal injection of MC-LR is 50 µg/kg body weight (bw); whereas mice LD50s for other microcystin variants range from 50 to >1,200 µg/kg bw (Carmichael, 1997). Chronic *Microcystis* exposure to laboratory mice resulted in liver injury, increased incidence of pneumonia, decreased survival, reduced brain size of neonates, and skin tumor promotion (Falconer et al., 1988; Falconer, 1991).

Microcystins have been related to many accidental animal poisonings. In Japan, 20 spot-billed ducks (*Anas paecilorhyncha haringtoni*) died from acute exposure to microcystins in a pond that became eutrophic from an influx of untreated sewage (Matsunaga, 1999). In England, ingestion of water from a waste storage reservoir containing a bloom of *Microcystis* was linked to the death of 20 lambs from an adjacent farm and 15 neighborhood dogs (Falconer, 1991). In addition, algal toxins may also initiate avian botulism outbreaks (Murphy et. al., 2000).

Bacterial Pathogens

Pasteurella multocida, *Yersinia* spp., *Salmonella* spp., *Erysipelothrix* spp., *Clostridium botulinum*, and *Escherichia coli* are bacteria associated with animal waste. The U.S. Geological Survey (USGS) National Wildlife Health Center (NWHC) has

identified these organisms as known or suspected waterfowl pathogens (USGS, 1999a and 2001).

Pasteurella multocida is highly infectious and can cause avian cholera in waterfowl. Infections of *P. multocida* in waterfowl are usually acute, often resulting in death within 6 to 12 hours, but also can be carried latently by birds and result in disease only under conditions of animal stress (USGS, 1999a). Environmental endurance of *P. multocida* may contribute to the length of time a seasonal outbreak persists (Rosen and Bischoff, 1950; Price and Brand, 1984) but is unlikely a source of outbreaks from one year to the next. Many avian cholera outbreaks have historically occurred among migrating waterfowl populations in the RWB and have resulted in substantial waterfowl mortality events (Zinkl et al., 1977; Price and Brand, 1984).

Yersiniosis in animals is characterized by gastroenteritis and diarrhea. *Yersinia* spp. are known to have established pathogen potential in animals, and swine are an especially important reservoir (Aleksic and Bockemühl, 1999). The pathogenic potential of *Y. intermedia* has not been completely determined, especially in relation to wildlife (Aleksic and Bockemühl, 1999).

Salmonella spp. are divided into six subgroups and there are over 2000 different serotypes recognized (Popoff and Minor, 1997). The natural reservoir for salmonellae is the intestinal tract of warm-blooded and cold-blooded animals. The majority of infected animals are apparently sub-clinically ill animals that harbor and shed the pathogen. *Salmonella* can survive in the environment for up to nine months or more, increasing dissemination potential (Quinn et al., 1994). In wild birds, particularly songbirds, gulls, and terns, salmonellosis can cause massive mortality events (USGS, 1999a). Avian salmonellosis (*Salmonella typhimurium*) was first diagnosed as a major cause of avian disease within the Salton Sea ecosystem in 1989, which resulted in the death of an estimated 4,515 cattle egrets (*Bubulcus ibis*) (Friend, 2002).

Erysipelothrix spp. have widespread environmental distribution in soil and water. Although primarily considered a swine pathogen, the organism has been isolated from many mammalian, avian, and amphibian species (Quinn et al., 1994). The distribution of

Erysipelothrix is probably under-reported. The association of *Erysipelothrix* with wildlife and fish, including major mortality events in eared grebes (*Podiceps nigricollis*) at the Great Salt Lake (Jensen and Cotter, 1976) suggests its inclusion in pathogen screens for waterfowl exposed to swine wastewater (USGS, 2001).

Bacteria of the genus *Clostridium* cause more wild avian mortalities than any other pathogen (USGS, 1999a). Avian botulism is a food poisoning caused by the ingestion of type C toxins produced by the bacterium *Clostridium botulinum* (USGS, 1999a). *Clostridium botulinum* spores are resistant to environmental extremes (Smith et al., 1982) and are widely distributed in wetland sediments (Smith and Sugiyama, 1988 as cited in Rocke and Samuel, 1999); however, botulism outbreaks in birds is dependent on several ecological factors including optimal environmental conditions for spore germination and bacterial growth, suitable material or substrates that provide energy for bacterial replication, and a mechanism of toxin transfer to birds (USGS, 1999a). Wetlands that tend to have botulism outbreaks have a greater percent organic matter in sediment, negative redox potential in the water, water pH between 7.5 and 9, water temperature above 20 °C, and salinity below 2 ppt (Rocke and Samuel, 1999).

Escherichia coli often infect the respiratory tracts of birds, resulting in colibacillosis, a chronic respiratory disease (USGS, 1999a). Lesions common to colibacillosis include pericarditis (i.e., an inflammation of the transparent membrane that encloses the heart), and perihepatitis (i.e., an inflammation of the peritoneal covering of the liver) (USGS, 1999a). Acute mortality to young waterfowl from *E. coli* infections has been reported in unhygienic hatcheries (USGS, 1999a).

Antibiotics

Hastings Pork regularly administers three tetracycline antibiotics (tetracycline, chlortetracycline, and oxytetracycline), two macrolide antibiotics (lincomycin and tylosin) and one diterpene antibiotic (tiamulin) to protect swine from disease and promote growth (Owen Nelson, Hastings Pork Farm Manager, pers. comm., 1999).

Antibiotics have become environmental contaminants of concern as they are designed to be biologically active, are generally water soluble, and they often have a low biodegradability (Wollenberger et al., 2000). The environmental fate of tetracycline antibiotics and their concentrations in CAFO wastewater is largely unknown (Zhu et al., 2001). The few studies that have examined the potential adverse effects from trace levels of antibiotics in the environment have focused on human health concerns associated with antibiotic resistance (DuPont and Steele, 1987; Guardabassi et al., 1998; Goñi-Urriza et al., 2000). Although the benefits from growth-promoting properties of antibiotics in animal feed have been known since the late 1940s, there has been little research on the mechanisms of antibiotics and potential effects to wildlife from chronic exposure to antibiotics (DuPont and Steele, 1987; Halling-Sørensen et al., 1997).

Tetracycline antibiotics are widely distributed in the body and sequestered particularly in liver, kidney, bone, and dentine (EAEM, 1995). There is no evidence of tetracycline antibiotics having the potential to cause reproductive or developmental toxicity, or carcinogenic or genotoxic effects (EAEM, 1995). Tetracycline and oxytetracycline are not acutely toxic to the freshwater crustacean *Daphnia magna* at environmentally relevant concentrations (Wollenberger et al., 2000). However, chronic toxicity tests on reproduction with *D. magna* indicated 50 percent of the population exhibited decreased reproductive output at concentrations of 46.2 mg/L and 44.8 mg/L for oxytetracycline and tetracycline, respectively (Wollenberger et al., 2000). Domestic rams receiving 20 mg/kg bw oxytetracycline for 6 or 10 days exhibited a decrease in spermatozoa motility and live spermatozoa count; however, these effects discontinued within 60 days following cessation of the treatment (HSDB, 1998). Tetracycline administered to pregnant rats resulted in neonates with discolored lens, cornea and sclera (HSDB, 1998).

Tiamulin inhibits protein synthesis at the ribosomal level (EAEM, 1995). Tiamulin chronic toxicity testing with *D. magna* resulted in decreased reproductive output for 50 percent of the population at 5.4 mg/L (Wollenberger et al., 2000). An oral dose of 50 mg/kg bw tiamulin to rats resulted in decreased testosterone-stimulated growth

of seminal vesicles with a no observed effect level (NOEL) of 15 mg/kg bw; however, dose levels up to 20 times the NOEL did not produce effects on reproductive performance, fertility, mass of gonads, or pathology (EAEM, 1995). Another study focused on reproductive effects to rats indicated no effects on fertility, growth, and survival of offspring at an oral dose of 100 mg/kg bw/day for 71 days prior to mating (EAEM, 1995).

Human treatment with lincomycin causes diarrhea in approximately 10 percent of patients and colitis (i.e., inflammation of the colon) in 1 percent of patients (Kelly et al., 1994). The antibiotic not only works against pathogenic bacteria but also against bacteria that are part of the digestive-tract flora of a healthy individual. When not kept in check by beneficial bacteria, microorganisms such as clostridia have the opportunity to grow excessively and release toxins resulting in colitis (Silva and Fekety, 1981). Less frequent toxic effects from lincomycin exposure in humans include cardiac arrhythmias, dermatitis, nephrotoxicity, hepatotoxicity, and various hematological abnormalities (HSDB, 1998).

Hormones

Run-off from CAFO wastes has been identified as an important source of synthetic and natural hormones to aquatic ecosystems (Shore, 1995; Kolpin et al., 2002; Burnison et al., 2003; Soto et al., 2004). Unlike the cattle industry, the swine industry does not use synthetic hormones (USEPA, 2003). Although synthetic hormones can be more potent and persistent than natural hormones, natural hormones can still exert effects on wildlife at low concentrations (Schiewer et al., 2001; Atienzar et al., 2002; Gross et al., 2003) and also may be more prevalent in the environment (Desbrow et al., 1998). Natural hormones associated with swine waste include testosterone, 17- β estradiol (E_2), estrone (E_1), and the phytoestrogen equol.

Risk to wildlife exposed to elevated levels of natural hormones is largely unknown due to a lack of data on environmental transport and fate in different environmental media (Kolpin et al., 2002; Ying et al., 2002). In sewage-treatment plant

effluent, E₂ is quickly metabolized to the less estrogenic E₁ by bacteria in sewage sludge, and the risk of extensive accumulation of natural estrogens in the environment is believed to be small (Lee and Liu, 2002). However, natural estrogens originating from swine waste were only degraded up to 20 percent after 12 weeks of storage at 20-23 °C (Lange et al., 2002), indicating a potential for concentrations to increase, especially under chronic exposure scenarios.

Most research on hormone contamination of natural waters has focused on sewage-treatment plants and exposure to fish (Routledge et al., 1998; Harries et al., 1999; Rogers-Gray et al., 2000; Sole et al., 2000) and there is a lack of information on the potential exposure and effects to other species (Kolpin, 2002; Lange et al., 2002). There are no studies that have evaluated waterfowl or shorebird exposure to natural hormones from ingestion of contaminated water or food items. However, in human health risk assessment there is a concern that ingestion of natural hormone residues in meat could result in neurobiological, developmental, reproductive, immunological, mutagenic and carcinogenic effects (European Commission, 1999).

METHODS

Sample collections for this study were aimed at comparing CAFO contaminants in sediments, water, and biota from the canals and created wetlands, which receive wastewater from the lagoons, and McMurtrey Marsh, a site adjacent to Hastings Park that does not receive swine wastewater effluent. McMurtrey Marsh was often too dry to sample; therefore, data collected from other studies also were used for comparisons. A more detailed description of the methods for each analysis is provided below.

Algal Toxins

Service personnel from the Nebraska Ecological Services Field Office (NEFO) examined the created wetlands for cyanobacteria blooms on a total of 17 occasions during the spring, summer, and fall of 2000. Phytoplankton samples were collected and viewed under a light microscope to determine presence or absence of cyanobacteria. If the algal community appeared to be dominated by cyanobacteria with the potential to form toxic blooms (e.g. *Anabaena*, *Aphanizomenon*, *Nodularia*, *Nostoc*, *Oscillatoria*, and *Microcystis*), then a concentrated sample was collected using a 67 micro-meter (μm) phytoplankton tow net. Concentrated samples were placed in 125 milliliter (ml) brown plastic containers and immediately shipped on dry ice to the U.S. Geological Survey Biological Resource Division's Columbia Environmental Research Center (CERC). Concentrations of hepatotoxins (i.e., microcystins reported as total microcystin LR) were determined by enzyme linked immunosorbent assay (ELISA). Microcystin and nodularian toxin variants were identified, to further characterize toxicity, by high-pressure liquid chromatography (HPLC). A more detailed description of the ELISA and HPLC methods used to analyze water samples collected for this study is available in Echols (2001).

Antibiotics

In March, June, and October, 2000, Service personnel collected water and sediment samples for pharmaceuticals analysis from lagoons, canals, created wetlands, and McMurtrey Marsh. Water samples were placed in pre-cleaned 500 ml amber glass bottles and stored on ice or chilled until extraction. Sediment samples were stored in 160 ml glass jars and kept frozen. The University of Nebraska-Lincoln Water Sciences Laboratory performed a tetracycline scan and a macrolide scan to analyze water and sediment samples for tetracycline, oxytetracycline, chlortetracycline, tiamulin, lincomycin, and tylosin. Tetracycline antibiotic concentrations were determined by solid phase extraction followed by detection with liquid chromatography and ion-trap electrospray ionization mass spectrometry as described by Zhu et al. (2001). Macrolide antibiotics were measured using solid phase extraction and liquid chromatography-tandem mass spectrometry according to procedures developed by Snow et al. (2003).

Bacteria Pathogens

Sediment and water samples were tested for microbial pathogens likely to occur in swine (i.e., *Clostridium botulinum* type C, *Salmonella* spp., *Pasteurella multocida*, *Yersinia* spp, *Erysipelothrix* spp, fecal coliforms, and fecal streptococci). Personnel from the NWHC and NEFO collected samples in April, June, and October 2000, from 14 stations at the study site (Figure 4). Water samples were collected in 1 L sterilized bottles and top sediment (i.e., < 10 centimeters) was placed in 125 ml sterilized containers. Water samples were kept on ice and processed within 6 hours of collection. Sediment samples (~0.25 grams) were placed into microcentrifuge tubes and stored frozen until analyzed. All samples were analyzed at the NWHC. Fecal coliform and fecal streptococci counts were obtained by membrane filtration (Clesceri et al., 1998). Samples with fecal coliform or fecal streptococci colony forming units that were too numerous to count (i.e., the agar plates were too overgrown to distinguish colony units) were not included for statistical analysis. The presence of *P. multocida*, *Salmonella*,

Yersinia and *Erysipelothrix* was verified biochemically by either the API-20E or Vitek systems (bioMerieux, St. Louis, Missouri). The presence of Botulism type C spores was determined using a DNA isolation kit (UltraClean™, MoBio Laboratories, Inc., Solana Beach, CA). Although not part of the original study plan, *Salmonella* and *Escherichia coli* isolates were tested for antibiotic resistance using standardized materials. A more detailed description of the methods used to determine antibiotic resistance, pathogen counts, and isolates is provided by USGS (2001).




Figure 4. Locations of samples collected for identification of bacteria pathogens at Hastings Pork and McMurtrey NWR, Clay County, Nebraska, 2000.

Hormones

Determining concentrations of hormones in water was not part of the original study plan; however, funding allowed for a limited analysis. Water samples were collected in 50 ml Falcon tubes, kept on ice, and shipped to the USGS Florida Caribbean Science Center where they were analyzed directly (i.e., without preparing an extract) for 17- β estradiol and testosterone using ELISA methods.

Waterfowl Use

The Rainwater Basin Wetland Management District performed waterfowl brood surveys in and around the created wetlands to determine waterfowl use. Surveys were conducted in July and September of 2000 and July and August of 1999. The number of broods, species, age class, and number of ducklings were recorded.

Invertebrate Diversity

Wetland benthic invertebrates were sampled three times during the year in spring, summer, and fall. Invertebrates were sampled using a 20 centimeter (cm) diameter stovepipe sampling core. Water within the pipe was decanted through a 297 μ m sieve and sediments were collected until firm ground was reached. For each created wetland and McMurtrey Marsh, a composite sample from six locations on each wetland was preserved in 10 percent formalin. There was a total of nine samples, with each sample representing one of three seasons (spring, summer, fall) for one of three sites, McMurtrey Marsh, created wetland 1 (CW1) and created wetland 5 (CW5). Invertebrates were randomly sub-sampled following standard USEPA rapid assessment procedures (Barbour et al., 1999) by picking all individuals within four randomly selected 6 by 6 cm grids. Analysis of invertebrates focused on taxa richness as our methods did not allow for density estimates. EcoAnalysts, Inc. (Moscow, Idaho) identified (usually to genus) all invertebrates in each subsample.

Water Quality

Temperature, dissolved oxygen, conductivity, salinity (YSI model 85) and pH (Accument[®] AP61) measurements were taken every 2 weeks on all created wetlands and McMurtrey Marsh from March 14 to November 07, 2000. In addition, water samples were collected monthly from McMurtrey Marsh, CW1, and CW5 from March to October 2000, and analyzed for total Kjeldahl nitrogen, ammonia, nitrates, and total phosphorus (Servi Tech Laboratories, Hastings, Nebraska). Water quality from the created wetlands and McMurtrey Marsh were compared to water quality data collected by the NDEQ for 30 stations at 15 RWB wetlands (NDEQ, 1997).

Trace Elements

Water, sediment, and invertebrate samples were submitted to the Service's Patuxent Analytical Control Facility (PACF) for trace metal analysis (Table 1). Samples were collected in USEPA certified clean glass containers and sampling equipment was decontaminated between sites. Water samples were collected in 500 ml containers and 2 ml of nitric acid were added to each sample to obtain a pH near 2. Sediment samples were collected into 1,500 ml containers with a stainless steel spoon in all areas except the lagoons where samples were collected using a ponar dredge. Forceps were used to collect chironomids from sediments filtered by a 1.1 mm mesh size sieve. Care was taken to select chironomids of all sizes as chironomid size can alter metal uptake (Krantzberg, 1989).

Inductively coupled plasma atomic emission spectrometry was used to determine concentrations of Al, B, Ba, beryllium (Be), Cd, chromium (Cr), Cu, Fe, magnesium (Mg), Mn, Mo, Ni, Pb, strontium (Sr), V, and Zn. Mercury (Hg) concentrations were determined by cold vapor atomic absorption. Graphite furnace atomic absorption was used to measure As, Se, and small concentrations of Pb and Cd.

Concentrations of trace elements in sediment were compared among Hastings Pork lagoons, canals, created wetlands, and Rainwater Basin wetlands. Data from water and sediment samples collected at McMurtrey Marsh during this study were pooled with

data on trace element concentrations from other studies on Rainwater Basin wetlands to increase the sample size of the “control wetlands” for comparison with the created wetlands, canals and lagoons. The RWB wetland sediment data set included 10 samples collected in 1993 and 1994 from RWB wetlands at McMurtrey NWR, Harvard WPA, Massie WPA, Eckhart WPA, and Smith WPA (PACF catalog numbers 6050045 and 6050027). The RWB wetland water data set included 16 water samples collected by the Service in 1991 and 1993 (catalogs 6050017 and 6050045, respectively). Not all samples collected were analyzed for the same trace elements; therefore, sample sizes for RWB wetland sediment and water data sets varied from 2 to 18 (Appendix, Table A.1).

Data on trace element concentrations in chironomids from wetlands in Nebraska is lacking; therefore, trace element concentrations in chironomids collected from the Hastings Pork created wetlands and canals were compared to those in chironomids collected at two National Irrigation and Water Quality Program (NIWQP) sites, namely, the Stillwater NWR in west-central NV (Tuttle et al., 1996) and the Sun River Irrigation Project Area in west-central Montana (NIWQP, 2003). Data on trace element concentrations in chironomids from these two areas were obtained through PACF and included catalogs 1070005, and 1070009 to 1070012 for Stillwater NWR and 6070002, 6070006, 6070010, 6070019, 6070029, 6070030, 6070033, 6070041, 6070042, 6070044, 6070049, 6070051 for sites within the Sun River Irrigation Project. Concentrations of Al, As, B, Hg, and Zn in biota, sediment and water samples collected from Stillwater NWR frequently exceeded concentrations associated with adverse biological effects (Tuttle et al., 1996); therefore, Stillwater NWR data represented a contaminated site. The Sun River Irrigation Project Area contained concentrations of Se in biota, sediment and water samples above toxicity thresholds (Palawski et al., 1991), but most other trace elements were typically below levels of concern (Bill Olsen, U.S. Fish and Wildlife Service Contaminants Specialist, pers. comm., 2003). Therefore, with the exception of Se, concentrations of trace elements in chironomids from this site represented an uncontaminated site.

Statistical Analyses

Statistical calculations were performed in software from the Statistical Analysis System (SAS) Institute (either JMP[®] Version 5 or SAS[®] Version 8.2). Data were typically nonparametric; therefore, a Kruskal-Wallis nonparametric one-way analysis of variance was used to test significance among three groups and a Wilcoxon rank sums test was used to test significance between groups. Differences among three or more groups were analyzed using PROC MIXED in SAS (SAS Institute, 2001). If more than 50 percent of the samples analyzed were above the detection limit for a particular trace element, then half the detection limit was used in place of those below the detection limit for statistical analyses, unless otherwise noted. Trace elements below the detection limit for more than 50 percent of the samples were not analyzed statistically.

Table 1. Summary of samples collected for trace element analyses at Hastings Pork and McMurtrey Marsh, 2000 to 2001.

| Date Collected | PACF Catalog ID | Site | Laboratory | Number of Samples per Matrix | | |
|----------------|-----------------|--------|------------|------------------------------|-------|---------------|
| | | | | Sediments | Water | Invertebrates |
| March, 2000 | 6050062 | CW1 | RTI | 1 | 1 | 0 |
| | | CW5 | | 1 | 1 | 0 |
| | | Canal | | 1 | 1 | 0 |
| | | MM | | 1 | 1 | 0 |
| June, 2000 | 6050065 | CW1 | MRI | 1 | 1 | 2 |
| | | CW5 | | 1 | 1 | 2 |
| | | Canal | | 1 | 1 | 0 |
| October, 2000 | 6050066 | CW1 | RTI | 1 | 1 | 1 |
| | | CW5 | | 1 | 1 | 1 |
| | | Canal | | 1 | 1 | 0 |
| | | MM | | 1 | 1 | 0 |
| July, 2001 | 6050091 | CW1 | GERG | 2 | 0 | 0 |
| | | CW2 | | 2 | 0 | 1 |
| | | CW3 | | 2 | 0 | 1 |
| | | CW4 | | 2 | 0 | 1 |
| | | CW5 | | 2 | 0 | 1 |
| | | CW6 | | 2 | 0 | 1 |
| | | CW7 | | 2 | 0 | 1 |
| | | Lagoon | | 10 | 0 | 0 |
| | | Canal | | 4 | 0 | 2 |
| | | MM | | 0 | 0 | 0 |
| TOTAL | | | | 39 | 11 | 14 |

Note: CW = created wetland; MM = McMurtrey Marsh; PACF = Patuxent Analytical Control Facility, Laurel, MD; ID = identification number; Laboratory = the laboratory that performed the analytical analysis (i.e., RTI = Research Triangle Institute, MRI = Midwest Research Institute, GREG = Geochemical & Environmental Research Group, Texas A&M).

RESULTS

Algal Toxins

Cyanobacteria genera (i.e., *Microcystis*, *Anabaena*, *Hapalosiphon*, *Nodularia*, and *Oscillatoria*) were identified in four of seven created wetlands sampled at Hastings Park in 2000 (created wetlands 3, 4, 6, and 7). Algal blooms dominated by *Microcystis* were observed in three created wetlands (CW 4, 6, and 7) on October 2 and 18, 2000. Six samples were collected from these blooms. All six samples contained microcystins reported as total microcystin-LR. The mean total microcystin-LR concentration determined by ELISA methods was 81 nanograms (ng) per mg (standard error = 24 ng/mg). Analysis by HPLC indicated microcystin-RR was the dominant microcystin variant in all six samples with microcystin-LR detected at concentrations ranging from 0.1 to 1.6 ng/mg (Table 2).

Table 2. Total microcystin (MC) concentrations in algal blooms from the Hastings Park created wetlands, Clay County, Nebraska, 2000.

| Sample ID and date collected | ELISA Total MC-LR | HPLC MC-LR | HPLC MC-RR | HPLC MC-YR | HPLC MC-LA |
|------------------------------|-------------------|------------|------------|------------|------------|
| CW4 10/2/2000 | 200 | ND | 120 | ND | ND |
| CW6 10/2/2000 | 60 | 0.1 | 0.6 | ND | ND |
| CW7 10/2/2000 | 73 | 0.3 | 19 | ND | ND |
| CW4 10/18/2000 | 61 | ND | 22 | ND | ND |
| CW6 10/18/2000 | 50 | 1.6 | 4.9 | ND | ND |
| CW7 10/18/2000 | 41 | ND | 4.2 | ND | ND |

Note: all microcystin concentrations are in ng/mg dry weight. ND=non-detect.

Antibiotics

The total number of water samples (n) collected for determining concentrations of antibiotics in water was 32 and included collections from lagoons (n=14), canals (n=10), created wetlands (n=6), and McMurtrey Marsh (n=2). In addition, nine sediment samples were collected in October from the lagoons (n=3), canals (n=3), created wetlands (n=2), and McMurtrey Marsh (n=1). Tetracycline, oxytetracycline, chlortetracycline, tiamulin, lincomycin, and tylosin were detected in either water or sediment samples collected from Hastings Pork lagoons and canals. In the created wetlands, only oxytetracycline was detected in sediments (41 ng/g); however, sample size for created wetlands was small (n = 2 and 6 for sediments and water, respectively). No antibiotics were detected in McMurtrey Marsh.

Tetracycline, oxytetracycline, chlortetracycline and tiamulin were frequently detected in sediments or water from the Hastings Pork lagoons and canals (Appendix, Table A.7). Oxytetracycline was detected in all water samples collected from the lagoons and in all sediment samples collected from the lagoons and canals. Average concentrations of oxytetracycline were more than 60 times greater in sediments than water (Figures 5 and 6). Tiamulin was detected in all water samples collected from the lagoons and in 2 of 3 sediment samples from the lagoons. In the Hastings Pork canals, tiamulin was detected in 80 percent and 30 percent of the water samples and sediment samples collected, respectively. Chlortetracycline was detected in all 6 sediment samples from lagoons and canals. Concentrations of chlortetracycline in sediments ranged from 311 to 6,430 ng/g in the lagoons and from 108 to 1,800 ng/g in the canals. Tetracycline was detected in 5 of 6 sediment samples and concentrations ranged from 119 to 1,328 ng/g in the lagoons and from 50 to 98 ng/g in the canals. Tetracycline and chlortetracycline in water samples were generally below the detection limit of 10 µg/L. In lagoon water samples (n=11), lincomycin was only detected twice (780 µg/L and 53 µg/L) and tylosin was only detected once (11.4 µg/L). Lincomycin and tylosin were not detected in lagoon sediments (n = 6).




Figure 5. Mean (\pm S.E.) concentrations of antibiotics in sediments from lagoons and canals at Hastings Park, Clay County, Nebraska, 2000. The number of detects per number of samples analyzed is displayed above each standard error bar. Note: tiamulin was detected once in canal sediments at 29 ng/g (data not shown).




Figure 6. Mean (\pm S.E.) concentrations of antibiotics in water from lagoons and canals at Hastings Park, Clay County, Nebraska, 2000. The number of detects per number of samples analyzed is displayed above each standard error bar.

Bacterial Pathogens

Bacterial results indicated the occurrence of fecal coliforms, fecal streptococci, *Salmonella* spp., *Yersinia* spp., and *Clostridium botulinum* type C in the created wetlands, lagoons and canals. Fecal coliform and fecal streptococci colony forming units (cfu) were counted in samples collected from the lagoons (n = 12), canals (n=12), created wetlands (n=6), a drainage ditch leading into McMurtrey NWR (n=3), and McMurtrey Marsh (n=6)(See Figure 4 for site locations). Fecal streptococci and fecal coliform colonies were too overgrown to distinguish colonies (too numerous to count) on 12 occasions from samples collected at the created wetlands, canals, and lagoons (Appendix, Table A.2). Annual mean counts of fecal coliform and fecal streptococci were highest in the lagoons and lowest in McMurtrey Marsh and the drainage ditch (Table 3). Fecal coliform and streptococci cfu per 100 ml (cfu/100ml) often varied considerably within sites and by season (Figure 7). Although this variation precluded significant differences in mean counts of cfu/100ml between many of the sites, the created wetlands had a significantly ($P < 0.05$) greater mean number of fecal streptococci cfu/100 ml when compared to McMurtrey Marsh (Table 3).

Table 3. Mean annual fecal coliforms and fecal streptococci colony forming units per 100 ml of water from sites at Hastings Pork and McMurtrey Marsh, Clay County, Nebraska, 2000.

| Pathogen | Site | n | Mean \pm S.E. | min | max | Significance* |
|--------------------|--------|----|----------------------|-------|---------|---------------|
| Fecal Coliforms | Lagoon | 9 | 164,894 \pm 81,990 | 4,000 | 770,000 | A |
| | Canal | 12 | 10,145 \pm 7,297 | 100 | 90,000 | B |
| | CW | 6 | 1,277 \pm 523 | 100 | 3,340 | B |
| | Ditch | 2 | 247 \pm 168 | 79 | 415 | B |
| | MM | 6 | 1,208 \pm 495 | 50 | 2,650 | B |
| Fecal Streptococci | Lagoon | 10 | 46,280 \pm 15,452 | 3,250 | 143,500 | A |
| | Canal | 11 | 4,768 \pm 1,067 | 600 | 12,100 | B |
| | CW | 5 | 27,472 \pm 24,432 | 200 | 125,000 | AB |
| | Ditch | 2 | 147 \pm 112 | 35 | 258 | BC |
| | MM | 6 | 82 \pm 46 | 5 | 300 | C |

*Different letters indicate significance ($p < 0.05$) as determined by a Kruskal-Wallis test followed by pairwise Wilcoxon rank sums tests. CW = created wetland, MM = McMurtrey Marsh, n = sample size and does not include samples in which fecal coliforms or streptococci were too overgrown to distinguish colonies (see Appendix table A. 2).




Figure 7. Mean (\pm S.E.) fecal coliform and fecal streptococci counts for water samples collected from created wetlands (CW), canals, lagoons, and McMurtrey Marsh (MM) in Clay County, Nebraska, 2000. N = the number of samples analyzed for each site per season; however, samples were not collected from MM in June of 2000 due to dry conditions.

Sixteen isolates of *Salmonella* spp. and 24 isolates of *Yersinia* spp. were recovered during this study (Appendix, Tables A.3 and A.4). All *Salmonella* spp. isolates were recovered from water samples, no isolates were obtained from sediments. *Salmonella* isolates were recovered from the Hastings Pork lagoons, canals, created wetlands, and a ditch leading into McMurtrey NWR, but were not recovered from McMurtrey Marsh (Figure 8). *Salmonella* serotypes isolated included Newport, Infantis, Muenchen, and Typhimurium (Copenhagen) (Appendix, Tables A.3 and A.4). *Yersinia* spp. isolates were recovered from both water and sediment samples with water being the primary source. The most common isolate was *Yersinia intermedia*, accounting for 14 of the 24 isolates. *Yersinia enterocolitica* accounted for 4 of 24 isolates.

Bacterial resistance to multiple antibiotics was detected for *Salmonella* spp and *E. coli* isolates. Two of 16 *Salmonella* spp. isolates exhibited resistance to multiple antibiotics. Out of 31 *E. coli* isolates tested; 28 were resistant to at least one antibiotic, 12 were resistant to two or more antibiotics, 9 were resistant to four or more antibiotics and 1 was resistant to eight antibiotics tested. Resistance to tetracycline was most common for both *Salmonella* spp. (20 percent of isolates tested) and *E coli* (> 90 percent tested). A more detailed description of the antibiotic resistance results are provided by USGS (2001).




Figure 8. Number of isolates of *Yersenia* and *Salmonella* species from McMurtrey Marsh and Hastings Pork created wetlands, canals and lagoons, Clay County, Nebraska, 2000. N = the total number of samples analyzed for each site.

Hormones

Concentrations of testosterone and E₂ in water from the created wetlands, lagoons and canals (n = 4 total) were greater than detected at McMurtrey Marsh (n=1) and were similar to concentrations from contaminated sites as reported by others (Table 4). 17 beta estradiol was detected in 4 out of 5 samples and concentrations ranged from 49 picograms per milliter (pg/ml) at Created Wetland 1 to below detection limits (< 5 pg/ml) at McMurtrey Marsh. Testosterone was detected in all 5 samples and concentrations ranged from 206 pg/ml in the canal to 23 pg/ml at McMurtrey Marsh.

Table 4. Concentrations of 17- β estradiol and testosterone in water samples collected from McMurtrey Marsh and Hastings Pork created wetlands, canals and lagoons in Clay County, Nebraska, 2000, compared to those published by other studies.

| Hormone | Site | Measure | Con.(ppt) | Units | Method | Suspected Source | Citation |
|-----------------------|-------------------|------------|-----------|-------|----------------|--------------------------------|----------------------------|
| 17- β estradiol | Lagoon | N=1 | 21 | pg/ml | RIA | Swine CAFO | This Study |
| | Created Wetland 5 | N=1 | 7 | pg/ml | RIA | Swine CAFO | This Study |
| | Canal | N=1 | 31 | pg/ml | RIA | Swine CAFO | This Study |
| | Created Wetland 1 | N=1 | 45 | pg/ml | RIA | Swine CAFO | This Study |
| | McMurtrey Marsh | N=1 | 0 | pg/ml | RIA | Uncontaminated | This Study |
| | British Rivers | Max | 50 | ng/L | Yeast bioassay | Sewage | Desbrow et al., 1998 |
| | Farm pond | Max | 7.4 | ng/L | RIA | Cattle Farm | Irwin et al., 2001 |
| | Grassland run-off | Range | 50 to 150 | ng/L | ELISA | Poultry farm | Finlay-Moore et al., 2000 |
| | U.S. Streams | Median* | 9 | ng/L | GC/MS | Industry, sewage, agricultural | Kolpin et al., 2002 |
| | U.S. Streams | Max (n=70) | 93 | ng/L | GC/MS | Industry, sewage, agricultural | Kolpin et al., 2002 |
| Testosterone | Lagoon | N=1 | 131 | pg/ml | RIA | Swine CAFO | This Study |
| | Created Wetland 5 | N=1 | 76 | pg/ml | RIA | Swine CAFO | This Study |
| | Canal | N=1 | 206 | pg/ml | RIA | Swine CAFO | This Study |
| | Created Wetland 1 | N=1 | 176 | pg/ml | RIA | Swine CAFO | This Study |
| | McMurtrey Marsh | N=1 | 23 | pg/ml | RIA | Uncontaminated | This Study |
| | Grassland run-off | Range | 15 to 125 | ng/L | ELISA | Poultry farm | Finlay-Moore et al., 2000. |
| | U.S. Streams | Median* | 116 | ng/L | GC/MS | Industry, sewage, agricultural | Kolpin et al., 2002 |
| | U.S. Streams | Max (n=70) | 214 | ng/L | GC/MS | Industry, sewage, agricultural | Kolpin et al., 2002 |

Note: Con. (ppt) = Concentration in parts per trillion; GC/MS = gas chromatography/mass spectroscopy. Full citations provided in the References section.

Waterfowl Use

Hastings Park and Service personnel observed heavy use of the created wetlands by waterfowl and shorebird species in 1999, 2000, and 2001. Species observed included: mallard, green-winged teal (*Anas crecca*), blue-winged teal (*Anas discors*), wood duck (*Aix sponsa*), redhead (*Aythya americana*), gadwall (*Anas strepera*), scaups (*Aythya* spp.), bufflehead (*Bucephala albeola*), American wigeon (*Anas americana*), northern shoveler (*Anas clypeata*), northern pintail, ruddy duck (*Oxyura jamaicensis*), Wilson's phalarope (*Phalaropus tricolor*), spotted sandpiper (*Actitis macularia*), dowitchers (*Limnodromus* spp.), and killdeer (*Charadrius vociferus*). Broods of mallard, gadwall, wood duck, redhead, and teal were observed in surveys performed in August of 1999, and July and September of 2000, with mallards being most frequently observed. Waterfowl also were observed loafing in the lagoons and canals, although these areas appeared to be utilized less often and by fewer numbers compared to the created wetlands.

Invertebrate Analysis

Invertebrate abundance and diversity were markedly different between McMurtrey Marsh and created wetland sites evaluated (Table 5). Invertebrates of the order Trichoptera, Odonata, Copepoda, Arhynchobdellida and members of the class Ostracoda and Oligochaeta were present at McMurtrey Marsh and absent from the created wetlands. Benthic invertebrate communities in the created wetlands were dominated by Chironomidae (93 to 100 percent) during all three seasons. In comparison, chironomids made up 0 to 15 percent of the benthic invertebrate community at McMurtrey Marsh. The invertebrate community at McMurtrey Marsh appeared to be comprised mainly of Cladocera in the spring and summer and Odonata in the fall. Taxa diversity was greatest at McMurtrey Marsh during the spring, summer, and fall when compared to the created wetland sites (Figure 9). Chironomids of the genus *Tanypus* was the dominant taxa in created wetlands for all seasons sampled with the exception of CW1 in the spring when the chironomid genus *Glyptotendipes* was dominant.

Table 5. Total number of individuals by invertebrate order found in the spring, summer, and fall of 2000 from sites at McMurtrey Marsh and created wetlands 1 and 5.

| Order-Family or (Class) | Genus | McMurtrey NWR | | | Created Wetland 1 | | | Created Wetland 5 | | |
|-------------------------|-----------------------------|---------------|--------|------|-------------------|--------|------|-------------------|--------|------|
| | | Spring | Summer | Fall | Spring | Summer | Fall | Spring | Summer | Fall |
| Odonata | Coenagrion and Enallagma | - | - | 22 | - | - | - | - | - | - |
| | Libellulidae | - | - | 2 | - | - | - | - | - | - |
| Hemiptera | Buenoa sp. | - | - | - | - | - | - | - | - | 15 |
| | Corisella sp. | - | 1 | 2 | - | - | 4 | 1- | 6 | 32 |
| | Sigara sp. | - | - | - | - | - | - | - | 5 | - |
| Coleoptera | Berosus sp. | - | - | 2 | - | - | - | - | - | - |
| | Coleoptera | - | - | - | - | - | 2 | - | - | - |
| | Ilybius sp. | 1 | - | - | - | - | - | - | - | - |
| Trichoptera | Oecetis sp. | - | - | 1 | - | - | - | - | - | - |
| Diptera | Ceratopogon sp. | - | - | 7 | - | - | - | 7 | 7 | 6 |
| | Mallochohelea sp. | - | - | - | - | - | - | 1 | - | - |
| | Muscidae | - | - | - | - | - | 2 | - | - | 2 |
| | Sphaeromias sp. | - | - | 1 | - | - | - | - | 7 | - |
| 39 Diptera-Chironomidae | Chironomus sp. | - | - | - | 2 | 4 | 41 | 25 | 47 | 142 |
| | Clinotanypus sp. | - | - | 4 | - | - | - | - | - | - |
| | Glyptotendipes sp. | - | - | - | 104 | 20 | 4 | 58 | 53 | 133 |
| | Tanypus sp. | - | - | 5 | 24 | 646 | 373 | 542 | 328 | 499 |
| Arhynchobdellida | Erpobdellidae | 2 | 4 | 2 | - | - | - | - | - | - |
| Lumbricina | Oligochaeta | 10 | 8 | 6 | - | - | - | - | - | - |
| Cladocera | | 21 | 21 | - | - | 6 | - | - | - | - |
| Ostracoda (Class) | | 2 | 1 | - | - | - | - | - | - | - |
| Copepoda (Class) | Calanoida | 8 | 1 | 4 | - | - | - | - | - | - |
| | Total Number of Individuals | 44 | 36 | 58 | 130 | 676 | 426 | 643 | 453 | 829 |
| | Percent Chironomids | 0% | 0% | 16% | 100% | 99% | 98% | 97% | 94% | 93% |




Figure 9. Shannon-Weaver taxa diversity for benthic invertebrates collected in the spring, summer, and fall from McMurtrey Marsh, Created Wetland 1, and Created Wetland 5 in Clay County, Nebraska, 2000.

Water Quality

Dissolved oxygen, specific conductivity, and pH were significantly greater ($p > 0.05$) in created wetlands compared to McMurtrey Marsh and the RWB wetlands sampled by NDEQ (Figure 10). Salinity also was significantly greater ($p < 0.05$) in created wetlands (mean = 0.72 ± 0.04) compared to McMurtrey Marsh (mean = 0.13 ± 0.02). There was no significant difference in water temperature between McMurtrey Marsh (mean = 16.6 ± 2.5) and the created wetlands (mean = 17.2 ± 1.0). The created wetlands had significantly greater concentrations ($p < 0.05$) of total phosphorous, ammonia, and total nitrogen when compared to McMurtrey Marsh and RWB wetlands (Figure 11).




Figure 10. Mean (\pm SE) measurements of pH, dissolved oxygen, and specific conductivity in the Hastings Pork created wetlands, McMurtrey Marsh, and Rainwater Basin (RWB) wetlands in Clay County, Nebraska, 2000. Measurements were generally taken between 10:00 and 14:00. The sample size is given above each standard error bar. Different letters indicate significance ($p < 0.05$) as determined by a Kruskal-Wallis test followed by pairwise Wilcoxon rank sums tests.




Figure 11. Mean (\pm SE) concentrations of ammonia, total Kjeldahl nitrogen, and total phosphorus in created wetlands, McMurtrey Marsh, and Rainwater Basin wetlands in Clay County, Nebraska, 2000. The sample size is given above each standard error bar. Different letters indicate significance ($p < 0.05$) as determined by a Kruskal-Wallis test followed by pairwise Wilcoxon rank sums tests.

Trace Elements

Sediment. Concentrations of Cd, Cr, Cu, Mg, Mn, Ni, Se and Zn in sediment were significantly ($p < 0.05$) greater in the lagoons compared to the canals, created wetlands, and RWB wetlands (Table 6). In addition, Mo was only found in samples collected from lagoons, where it was detected in 8 of 10 samples (Appendix, Table A.8). Created wetlands had significantly ($p < 0.05$) greater concentrations of Al, B, Be, Cr, Mg, Mn, Sr, and V when compared to RWB wetlands (Table 6); whereas, RWB wetlands had significantly ($p < 0.05$) greater concentrations of Pb than lagoons, canals, and created wetlands (Figure 12). However, mean trace element concentrations in created wetlands and RWB wetlands were similar or less than background concentrations and did not exceed sediment quality guidelines or literature established toxicity thresholds (Table 7). Sediment toxicity thresholds and quality guidelines/benchmarks were exceeded in lagoon sediments for Cd, Cu, Ni, Mn, Se, and Zn. Trace element concentrations in canal sediments exceeded quality guidelines for Cu and Zn.

Water. Concentrations of trace elements in water from the created wetlands and canals were generally similar to those at McMurtrey Marsh; however, only 2 to 3 samples were analyzed for each site (Table 8). When data were pooled for the canals and created wetlands, these sites had significantly greater concentrations of B, Mg, and Mn compared to pooled data from the RWB sites (Figure 13). Concentrations of Se in water were detected in 3 of 9 samples from the Hastings Pork canals and created wetlands but Se was not detected in two water samples from McMurtrey Marsh. However, the detection limits for Se and Cd in water samples collected for this study exceeded their water quality criteria and/or their effects thresholds (Table 8). Concentrations of Zn in water samples from all sites generally exceeded the 0.03 mg/L “level of concern” at which toxic effects may occur in sensitive phytoplankton and invertebrate species (Suter and Tsao, 1996; USDI, 1998). Water quality criteria for Nebraska wetlands were exceeded for Al, Cd, Cu, Fe, and Pb in the created wetlands, canals, and McMurtrey Marsh and for As and Ni in the created wetlands (Table 8). Concentrations of Cu in water from the canals and

created wetlands exceeded Nebraska Cu aquatic life water quality criteria in all nine samples; whereas, samples from RWB wetlands exceeded Cu criteria concentrations in 2 of 11 samples (Table 8 and Appendix, Table A.5)

Invertebrates. Concentrations of trace elements in invertebrates from the created wetlands and canals, with the exception of Se, did not exceed any known toxicity thresholds for invertebrates or avian dietary items. Chironomids from the canals and created wetlands had concentrations of Se that were typically within the normal background (i.e., 0.4 to 4.5 mg/kg dw) for aquatic invertebrates (USDI, 1998). However, reproductive impairment in avian species can result from diets containing 3 to 8 mg/kg (USDI, 1998). Chironomids from the created wetlands exceeded 3 mg/kg Se in 4 of 10 samples analyzed. Concentrations of trace elements in chironomids from Stillwater NWR were generally significantly greater than those from Hastings Park, with the exception of Cr, Ni, Se and Zn (Table 9). Chironomids from the Sun River Irrigation Project Area had significantly lower concentrations of Al, Ba, Cr, Cu, Fe, Mn, Pb, and V compared to chironomids from the created wetlands and significantly greater concentrations of Se compared to chironomids from Stillwater NWR or the created wetlands (Table 9).

Table 6. Results from Proc Mixed in the Statistical Analysis System (SAS) for trace elements in sediments from Rainwater Basin wetlands and Hastings Pork lagoons, canals, and created wetlands in Clay County, Nebraska.

| Trace Elements | p values | Results from PROC MIXED in SAS | | | |
|----------------|----------|--------------------------------|-----------------------|----------------------|----------------------|
| Al | <0.0050 | CW ^A | Canals ^B | Lagoons ^B | RWB ^B |
| B | <0.0001 | CW ^A | Lagoons ^{AB} | Canals ^{BC} | RWB ^C |
| Be | <0.0024 | CW ^A | Lagoons ^A | Canals ^{AB} | RWB ^B |
| Cd | <0.0001 | Lagoons ^A | Canals ^B | RWB ^B | CW ^B |
| Cr | <0.0001 | Lagoons ^A | CW ^B | Canals ^{BC} | RWB ^{BC} |
| Cu | <0.0001 | Lagoons ^A | Canals ^B | CW ^C | RWB ^C |
| Mg | <0.0001 | Lagoons ^A | Canals ^B | CW ^B | RWB ^C |
| Mn | <0.0001 | Lagoons ^A | CW ^B | Canals ^{BC} | RWB ^C |
| Ni | <0.0001 | Lagoons ^A | Canals ^B | CW ^{BC} | RWB ^C |
| Pb | <0.0001 | RWB ^A | CW ^B | Canals ^C | Lagoons ^D |
| Se | <0.0001 | Lagoons ^A | Canals ^B | BDL | BDL |
| Sr | <0.0001 | Lagoons ^A | Canals ^{AB} | CW ^B | RWB ^C |
| V | <0.0001 | Lagoons ^A | CW ^A | Canals ^B | RWB ^C |
| Zn | <0.0001 | Lagoons ^A | Canals ^B | CW ^C | RWB ^C |

Note: CW = created wetlands, RWB = Rainwater Basin Wetlands. Different superscript letters indicate significant differences among sites. Sites are listed from left to right in decreasing order of mean concentration for each trace element. Trace elements tested that were not statistically different among sites include As, Ba, and Fe. Mo was only detected in lagoons. BDL = over 50 percent of samples tested were below the detection limit.




Figure 12. Mean (\pm SE) concentrations of lead (Pb) in sediments from Rainwater Basin wetlands and Hastings Pork created wetlands, canals, and lagoons, Clay County, Nebraska, 2000. The number of samples analyzed is displayed above each standard error bar. Different letters indicate significance ($p < 0.05$) as determined by “proc mixed” in SAS®

Table 7. Mean (\pm SE) trace element concentrations in sediments from the Hastings Park created wetlands and Rainwater Basin Wetlands compared to western background concentrations and effects thresholds.

| Trace Element | Lagoons (n=20) | Canals (n=7) | Created Wetlands (n=20) | Rainwater Basin Wetlands ¹ (n=12) | Western U.S. Background ² | Effects Thresholds |
|---------------|------------------------|-----------------------|-------------------------|--|--------------------------------------|--|
| Al | 10,690 \pm 1551 | 16,735 \pm 1167 | 17,267 \pm 1119 | 11,565 \pm 963 | 74,000 | 58,030 ^A |
| As | 4.8 \pm 0.5 | 4.3 \pm 0.3 | 4.5 \pm 0.3 | 3.6 \pm 0.3 | 7.0 | 9.79 ^B , 12.10 ^C |
| B | 13 \pm 1.1 | 8.9 \pm 2 | 15.9 \pm 1.3 | 5.2 \pm 1.3 | NA | No criterion |
| Ba | 195 \pm 5.9 | 220 \pm 23 | 215 \pm 15 | 202 \pm 23 | 670 | No criterion |
| Be | 1.08 \pm 0.07 | 0.85 \pm 0.11 | 1.09 \pm 0.08 | 0.64 \pm 0.01 | 0.97 | No criterion |
| Cd | 2.4 \pm 0.4 | 0.5 \pm 0.1 | 0.3 \pm 0.04 | 0.4 \pm 0.1 | NA | 0.59 ^C , 0.99 ^B |
| Cr | 42.6 \pm 2.5 | 21.3 \pm 3.5 | 21.3 \pm 1.5 | 14.6 \pm 1.0 | 56 | 43.4 ^B , 56.0 ^C |
| Cu | 325 \pm 50 | 39.3 \pm 9.3 | 16.3 \pm 1.1 | 15 \pm 1.0 | 27 | 31.6 ^B , 7.77 ^A , 270 ^D |
| Fe | 13,586 \pm 1884 | 14,291 \pm 752 | 17,101 \pm 892 | 15,099 \pm 2,154 | 26,000 | No criterion |
| Hg | <0.2 | <0.2 | <0.2 | 0 \pm 0 | NA | No criterion |
| Mg | 15,012 \pm 2280 | 6728 \pm 710 | 5773 \pm 321 | 2702 \pm 236 | NA | No criterion |
| Mn | 1025 \pm 134 | 436 \pm 102 | 450 \pm 49 | 311 \pm 40 | 480 | 819.0 ^E , 1,081 ^A , 1,673 ^C |
| Mo | 12 \pm 2.1 | BDL | BDL | BDL | 1.1 | No criterion |
| Ni | 26.8 \pm 1.6 | 18.6 \pm 1.2 | 17.4 \pm 0.9 | 14.4 \pm 1.2 | 19 | 22.7 ^B |
| Pb | 7.9 \pm 1.4 | 12.6 \pm 0.7 | 14.8 \pm 0.5 | 19.7 \pm 1.4 | 20 | 34.2 ^C , 35.8 ^B |
| Se | 6.1 \pm 0.8 | 1.4 \pm 0.4 | <1.0 | <1.0 | 0.34 | 4.0 ^E |
| Sr | 236 \pm 39 | 158 \pm 55 | 65 \pm 7.0 | 33.7 \pm 2.3 | NA | No criterion |
| V | 33.5 \pm 1.4 | 26.4 \pm 2.2 | 33.4 \pm 2.1 | 18.0 \pm 1.4 | 88 | No criterion |
| Zn | 2,134 \pm 407 | 189 \pm 45 | 72.1 \pm 5.8 | 60.3 \pm 2.8 | 65 | 121 ^B , 159 ^C , 1,532 ^A |

Note: SE = standard error, n = sample size, NA = not applicable, < = all samples were below the detection limit (value equals the greatest detection limit for all catalogs). Bold values indicate that the mean exceeded an effects threshold.

¹ USFWS, unpublished data (see Methods text for description)

² Background soil concentrations for the Western U.S. (Shakette and Boerngen, 1984).

^A = Probable Effects Concentration benchmark (Jones et al., 1997).

^B Sediment quality guideline threshold effects concentration below which harmful effects are unlikely to be observed (MacDonald et al., 2000).

^C = Toxic Effects Concentration benchmark below which effects are rarely expected to occur (Jones et al., 1997).

^D = No Effects Concentration benchmark (Jones et al., 1997).

^E = Toxicity threshold at which adverse effects on some fish and wildlife species may occur (USDI, 1998).

Table 8. Total recoverable concentrations (mg/L) of trace elements in water samples collected from McMurtrey Marsh and Hastings Pork canals and created wetlands, Clay County, Nebraska.

| Trace Element | Site and Dates (month/year) of Sample Collection | | | | | | | | | | | Effects Threshold (mg/L) |
|---------------|--|--------------|--------------|-------------------|--------------|--------------|-------------------|--------------|------------------|-----------------|---------------|---|
| | Canals | | | Created Wetland 1 | | | Created Wetland 5 | | | McMurtrey Marsh | | |
| | 3/00 | 6/00 | 10/00 | 3/00 | 6/00 | 10/00 | 3/00 | 6/00 | 10/00 | 3/00 | 10/00 | |
| Al | 0.321 | 0.879 | 0.154 | 3.610 | 2.640 | 0.173 | 14.700 | 8.710 | <0.111 | 0.657 | 11.900 | 0.087 ^A , 0.750 ^B |
| As | <0.0056 | 0.005 | 0.008 | 0.008 | 0.017 | 0.010 | 0.017 | 0.029 | 0.019 | <0.0056 | 0.012 | 0.016 ^A , 0.340 ^B |
| B | 0.37 | 0.55 | 0.41 | 0.25 | 0.55 | 0.43 | 0.20 | 0.71 | 0.38 | 0.05 | *0.111 | 6.0 ^C |
| Ba | 0.051 | 0.033 | 0.033 | 0.122 | 0.109 | 0.044 | 0.335 | 0.278 | 0.073 | 0.083 | 0.590 | 1.0 ^D |
| Be | <0.0011 | <0.0003 | <0.0006 | <0.0011 | <0.0003 | <0.0006 | <0.0011 | 0.001 | <0.0006 | <0.0011 | 0.002 | 0.005 ^A , 0.130 ^B |
| Cd | 0.0013 | <0.0004 | <0.0006 | <0.0011 | <0.0004 | <0.0006 | 0.0014 | <0.0004 | <0.0006 | <0.0011 | 0.0014 | *0.00025 ^A , *0.00590 ^B |
| Cr | <0.0056 | 0.003 | <0.0056 | <0.0056 | 0.003 | <0.0056 | 0.014 | 0.008 | <0.0056 | <0.0056 | 0.011 | *0.077 ^A , *0.592 ^B |
| Cu | 0.106 | 0.023 | 0.033 | 0.020 | 0.016 | 0.021 | 0.029 | 0.028 | 0.017 | 0.006 | 0.030 | *0.009 ^A , *0.013 ^B |
| Fe | 1.18 | 0.80 | 0.26 | 3.36 | 1.91 | 0.20 | 13.70 | 5.15 | 0.17 | 0.40 | 12.20 | 1.0 ^A |
| Hg | <0.0002 | <0.0002 | <0.0002 | <0.0002 | <0.0002 | <0.0002 | <0.0002 | <0.0002 | <0.0002 | <0.0002 | <0.0002 | 0.0005 ^A , 0.0014 ^B |
| Mg | 31.60 | 51.40 | 50.80 | 25.20 | 50.90 | 52.30 | 31.40 | 55.80 | 35.00 | 9.46 | 11.50 | 82.0 ^E |
| Mn | 0.31 | 0.11 | 0.10 | 0.29 | 0.24 | 0.09 | 0.87 | 0.56 | 0.13 | 0.01 | 0.85 | 1.0 ^A , 1.10 ^E |
| Mo | 0.017 | 0.005 | <0.056 | 0.007 | <0.004 | <0.056 | <0.006 | 0.014 | <0.056 | <0.006 | <0.056 | 10.0 ^F |
| Ni | 0.034 | 0.044 | 0.036 | 0.023 | 0.051 | 0.029 | 0.032 | 0.074 | 0.027 | <0.0056 | 0.017 | *0.052 ^A , *0.468 ^B |
| Pb | 0.009 | 0.001 | <0.0111 | 0.010 | 0.003 | <0.0111 | 0.025 | 0.009 | <0.0111 | <0.0056 | 0.045 | *0.0025 ^A , *0.0646 ^B |
| Se | <0.0056 | 0.004 | <0.0044 | <0.0056 | 0.003 | <0.0044 | <0.0056 | 0.007 | <0.0044 | <0.0056 | <0.0044 | 0.002 ^G , 0.005 ^A |
| Sr | 0.442 | 0.261 | 0.220 | 0.244 | 0.256 | 0.188 | 0.261 | 0.366 | 0.182 | 0.190 | 0.226 | 42.0 ^E |
| V | 0.006 | 0.017 | 0.006 | 0.017 | 0.019 | 0.009 | 0.046 | 0.040 | 0.012 | <0.0056 | 0.026 | 0.08 ^E |
| Zn | 0.566 | 0.101 | 0.103 | 0.066 | 0.058 | 0.040 | 0.094 | 0.067 | 0.013 | <0.0011 | 0.088 | 0.03 ^G , *117.00 ^B |

Note: > Indicates the sample was below the detection limit (value = detection limit); Date Col. = approximate date of sample collection; CW = created wetland; * indicates the criterion value based on a water hardness of 100 mg/L. Thresholds listed below are for dissolved concentrations unless stated otherwise. Bold numbers indicate that one or more of the following water quality criteria were exceeded:

^A = chronic aquatic life water quality criterion for wetlands (NDEQ, 2002).

^B = acute aquatic life water quality criterion for wetlands (NDEQ, 2002).

^C = level of concern threshold for aquatic invertebrates (USDI, 1998).

^D = Canadian Council of Ministers of the Environment (CCME) water quality guideline for the protection of aquatic life (CCME, 2002)

^E = The lowest chronic value benchmark above which toxic effects may occur in sensitive species (Suter and Tsao, 1996).

^F = concentration associated with frequent molybdenosis in sensitive species (Eisler, 1989).

^G = total recoverable Se toxicity threshold for apparent adverse effects to wildlife (USDI, 1998).




Figure 13. Mean (\pm SE) concentrations of boron (B), magnesium (Mg) and manganese (Mn) in water from Rainwater Basin wetlands and Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000. The number of samples analyzed is displayed above each standard error bar. Different letters indicate significance as determined by a Wilcoxon rank sums test.

Table 9. Results from Proc Mixed in the Statistical Analysis System (SAS) for trace elements in chironomids from Hastings Pork, Nebraska; Stillwater National Wildlife Refuge (NWR), Nevada; and the Sun River Irrigation Project Area, Montana.

| Trace Element | p values | Results of PROC MIXED in SAS and (mean ± standard error) | | |
|---------------|----------|--|--|---|
| Al | <0.0001 | Stillwater NWR ^A (9756 ± 768) | Hastings Pork ^B (5167 ± 525) | Sun River ^C (2285 ± 229) |
| As | <0.0001 | Stillwater NWR ^A (15.2 ± 0.9) | Sun River ^B (2.4 ± 0.2) | Hastings Pork ^C (1.5 ± 0.1) |
| B | <0.0001 | Stillwater NWR ^A (114.2 ± 9.2) | Sun River ^B (8.7 ± 0.9) | Hastings Pork ^B (7.7 ± 0.9) |
| Ba | <0.0001 | Stillwater NWR ^A (99 ± 7) | Hastings Pork ^B (64.7 ± 5) | Sun River ^C (36.2 ± 3) |
| Be | <0.0003 | Stillwater NWR ^A (0.41 ± 0.04) | Hastings Pork ^B (0.26 ± 0.04) | Sun River ^B (0.16 ± 0.03) |
| Cd | <0.0001 | Stillwater NWR ^A (1.13 ± 0.1) | Hastings Pork ^B (0.44 ± 0.1) | Sun River ^B (0.39 ± 0.1) |
| Cr | <0.0001 | Hastings Pork ^A (7.2 ± 0.7) | Stillwater NWR ^A (6.9 ± 0.5) | Sun River ^B (2.9 ± 0.3) |
| Cu | <0.0001 | Stillwater NWR ^A (30.0 ± 1.6) | Hastings Pork ^B (19.5 ± 1.7) | Sun River ^C (14.4 ± 0.6) |
| Fe | <0.0001 | Stillwater NWR ^A (10487 ± 677) | Hastings Pork ^B (4429 ± 463) | Sun River ^C (2682 ± 228) |
| Mg | <0.0001 | Stillwater NWR ^A (7951 ± 514) | Sun River ^B (3696 ± 290) | Hastings Pork ^C (2646 ± 158) |
| Mn | <0.0001 | Stillwater NWR ^A (295 ± 20.3) | Hastings Pork ^B (130 ± 12.9) | Sun River ^C (81 ± 7.5) |
| Ni | 0.0889 | Hastings Pork (10 ± 2) | Stillwater NWR (8 ± 1) | Sun River (6 ± 1) |
| Pb | <0.0001 | Stillwater NWR ^A (17.0 ± 2.1) | Hastings Pork ^B (5.0 ± 0.7) | Sun River ^B (3.0 ± 0.3) |
| Se | <0.0005 | Sun River ^A (10.4 ± 0.6) | Hastings Pork ^B (2.8 ± 0.4) | Stillwater NWR ^C (1.6 ± 0.2) |
| Sr | <0.0001 | Stillwater NWR ^A (206 ± 15) | Sun River ^B (41 ± 5) | Hastings Pork ^B (29 ± 4) |
| V | <0.0001 | Stillwater NWR ^A (34 ± 2) | Hastings Pork ^B (10 ± 1) | Sun River ^C (5 ± 0.5) |
| Zn | 0.1785 | Hastings Pork (91 ± 7) | Stillwater NWR (76 ± 4) | Sun River (76 ± 41) |

Note: Different superscript letters indicate significant differences among sites. Sites are listed from left to right in decreasing order of mean concentration (mg/kg dry weight) for each trace element. Mean Zn and Ni concentrations were not statistically different among sites.

DISCUSSION

Environmental pollution associated with CAFOs is a national issue (USDA and USEPA, 2003). Another national issue is the need to create and restore wetland habitat to remedy the perpetual decline of waterfowl and shorebird populations (North American Waterfowl Management Plan, 2003; Ducks Unlimited, 2003). The use of swine wastewater effluent from Hastings Pork to create waterfowl habitat is an attempt to simultaneously accommodate the need of CAFO managers to store and treat wastewater and the need for wildlife managers to provide habitat for waterfowl and shorebirds during the spring migration. However, wetland habitat created with swine wastewater may put these species at risk, if the water quality of these created wetlands is not suitable.

Although there are no known studies that specifically investigate risk to waterfowl exposed to swine wastewater, many of the constituents or environmental factors associated with swine waste are potentially harmful to waterfowl including Se (Heinz, 1996; USDI, 1998); cyanobacteria toxins (Matsunaga et al., 1999); *Salmonella* (Friend, 2002), eutrophication (Gaiser and Lang, 1998); and increased water conductivity (Mitcham and Wobeser, 1988; USDI, 1998). However, there also are many unknowns regarding swine waste exposure and effects to waterfowl and their habitat such as chronic exposure to antibiotics, natural hormones, and microcystin toxins.

The purpose of this study was to evaluate whether migratory waterfowl that utilize wetlands created from primary treated swine wastewater are likely exposed to contaminants including trace elements, salts, nutrients, cyanobacteria toxins, bacterial pathogens, and antibiotics. Waterfowl were chosen as the species of concern as the created wetlands were designed specifically to attract them. A CAFO contaminant exposure pathway from the lagoons, through the canals, and to the created wetlands was evaluated by collecting sediment, water, and invertebrate samples. In addition to the exposure assessment, habitat variables (water quality and invertebrate assemblages) were compared between the created wetlands and Federal wetlands managed for waterfowl habitat.

Results of this study indicate that many CAFO contaminants such as disease pathogens, nutrients, and some trace elements are passed through primary treatment lagoons and into the canals and created wetlands; whereas, many other trace elements and antibiotics appeared to be mainly trapped in lagoons. Trace element concentrations in sediment exceeded toxicity thresholds in lagoons and canals but not in created wetlands or RWB wetlands. Antibiotics also were frequently detected in lagoon sediments and water, but not in the created wetlands. When compared to McMurtrey Marsh and other RWB wetlands, created wetlands exhibited eutrophication; increased species richness of disease pathogens, pH, specific conductivity; and higher concentrations of nutrients and some trace elements.

Trace Elements

The comparison of trace element concentrations in sediment samples between RWB sites and Hastings Pork lagoons, identified B, Cd, Cr, Cu, Mg, Mn, Mo, Ni, Se, Sr, V, and Zn as CAFO contaminants. These trace elements are frequently detected in hog manure (Racz and Fitzgerald, 2001) and many of them (e.g., Cu, Cr, Se, and Zn) are supplied in feeds as nutritional supplements for disease suppression and growth promotion (Sims 1995 as cited by USEPA, 1998). The tendency for concentrations of trace elements to decrease in waterbodies down-gradient from the lagoons may indicate an effectiveness of the primary treatment in limiting their movement. However, concentrations of B, Cr, Cu, Mg, Mn, Ni, Se, Sr, V, and Zn in sediments and/or water from the created wetlands and canals appears to indicate that these trace elements are moving out of the lagoons. The transport of these trace elements to the created wetlands is likely due to the water solubility of the excreted trace element compounds. As much as 95 percent of dietary Cu in hog feed is subsequently excreted and much of it is in a readily soluble form (Payne et al., 1988). Boron compounds, especially from sewage and laundry products, also have high water solubility and conventional sewage treatment removes little or no boron (USEPA 1975 cited in Eisler, 1990).

Concentrations of Al, As, Cd, Cu, Fe, Ni, Pb, Se, and Zn in water from the created wetlands exceeded Nebraska aquatic life water quality criteria or literature established toxicity thresholds. However, the same criteria also were exceeded in McMurtrey Marsh for all these elements except As, Ni, and Se. In addition, the comparison of trace element concentrations detected in this study with Nebraska water quality criteria may not accurately evaluate whether wetland plant and invertebrate species are at risk. Recoverable metals were measured in this study, whereas Nebraska water quality criteria are generally based on measurements of dissolved metals. Water samples are filtered before an analysis for dissolved metals, whereas total recoverable analysis includes microorganisms and suspended particulates that are not filtered and thus result in higher concentrations. The total recoverable method was used in this study due to the highly productive nature of wetland systems where nutrients and toxins are quickly taken up by biota, leaving decreased concentrations in water (USDI, 1998). Concentrations of Al, Cd, and Fe in water were similar between created wetlands and McMurtrey Marsh and may reflect naturally high background concentrations (NDEQ, 1997). The greater concentrations of Pb in sediments and water from RWB wetlands compared to the created wetlands may reflect the presence of Pb shot from public hunting, even though steel or non-toxic shot has been required for all waterfowl hunting on Service RWB wetlands since 1980. Arsenic was detected at greater concentrations in water from the created wetlands compared to RWB wetlands indicating possible As contamination from swine waste. However, concentrations of As in sediments were similar across all sites and did not exceed sediment toxicity thresholds. Concentrations of Se in water from the created wetlands were generally below detection; however, sample sizes were small and the detection limits exceeded a 2 µg/L total recoverable Se toxicity threshold (USDI, 1998). Cu and Zn concentrations frequently exceeded Nebraska wetland water quality criteria; however, the high pH of the created wetlands likely limits their bioavailability and toxicity to wetland plants and invertebrates. Wetland macrophytes and algae were not analyzed for trace element concentrations; however, some wetland plant species can bioaccumulate copper to high concentrations (Buckley, 1994; Eisler, 1998a) and many

studies have reported trace element accumulation in soil and terrestrial plants exposed to applications of swine slurry (Christie and Beattie, 1989; Arzul and Maguer 1990).

Chironomids were selected as a potential food item for waterfowl and shorebirds as they appeared to be the most abundant benthic invertebrate at the created wetlands and are food items of major importance to blue-winged teal, northern pintail, mallard, gadwall, and redhead hens (Eldridge, 1990). Concentrations of trace elements in the chironomids did not exceed literature established toxicity thresholds and were generally significantly less than those found in chironomids from areas of known metal contamination. Trace element concentrations in chironomids at the created wetlands do not appear to represent a risk to waterfowl; however, when compared to concentrations in chironomids from Stillwater NWR (a contaminated site) or the Sun River Irrigation Project Area (a site with elevated Se), chironomids from the created wetlands appear to be accumulating some of the trace elements that were detected at high concentrations in lagoon sediments (e.g., Cr, Cu, Mg, Mn, Se, V, and Zn). Background concentrations of trace elements for chironomids from RWB wetlands are needed to better evaluate trace element bioaccumulation by chironomids in the created wetlands.

Antibiotics

The distribution of tetracycline antibiotics between sediment and water in the lagoon and canals is consistent with their high sorption coefficient in soil/sediment (K_d). The K_d for tetracycline and oxytetracycline can range from > 400 to 1,600 L/kg in soil (Tolls, 2001). The detection of tetracycline antibiotics in the canals suggests that even highly sorptive antibiotics are being transported away from the lagoons (Dr. Daniel Snow, Environmental Geochemist at the University of Nebraska Water Sciences Laboratory, pers. comm., 2004).

Antibiotic exposure to waterfowl and shorebirds at the created wetlands appeared to be essentially negligible; however, detection limits may have been too high. All six water samples collected from the created wetlands had oxytetracycline concentrations below the 10 µg/L detection limit. A national survey that consisted of 189 water samples

from 13 fish hatcheries reported that oxytetracycline was detected in 27 samples at a median concentration less than 0.05 mg/L and a one-time maximum concentration of 10 µg/L (Thurman et al., 2002). The same study concluded that the source of oxytetracycline was fish hatchery feed; therefore, the similar addition of oxytetracycline to hog feed would likely lead to its presence in the created wetlands.

There are no known studies that evaluate low-level chronic antibiotic exposure to avian wildlife; however, chronic toxicity tests on reproduction with *Daphnia magna* found 50 percent of the population exhibited decreased reproductive output at concentrations of 46.2 mg/L oxytetracycline (Wollenberger et al., 2000). Differences in wildlife sensitivity to environmental contaminants often occur between species and especially between classes (Calabrese and Baldwin, 1993). Avian specific toxicity evaluations would be needed to adequately evaluate potential adverse effects to migratory birds chronically exposed to low concentrations of oxytetracycline.

Bacterial Pathogens

Swine wastewater from Hastings Pork is apparently a source for some disease pathogens including fecal coliforms, fecal streptococci, *Salmonella* spp., and *Yersinia enterocolitica*; however, *Erysipelothrix* spp. and *P. multocida* were not detected. The lack of *Erysipelothrix* recoveries was unexpected because swine have been considered an important reservoir for this pathogen. Improvements to the *Erysipelothrix* recovery protocol may be needed to insure there are truly no *Erysipelothrix* spp. present in the study area (USGS, 2001). The absence of *P. multocida* in sediment and water samples collected for this study is not too surprising, as carrier animals are believed to be the most important reservoir and the bacteria are not believed to persist more than 2-4 weeks after carcasses are removed from a die-off event (Dr. Michael Samuel, Researcher at NWHC, pers. comm., 2004). There are no other known studies that have attempted to recover *P. multocida* from swine or cattle wastewater or sediments, although Smith (et al., 1989) found no association between the proximity of cattle feed lots and wetlands with frequent avian cholera outbreaks.

Avian cholera outbreaks at the created wetlands may still be a concern as disease transmission from carrier birds in the area has occurred historically and previous research is inconclusive regarding whether water quality characteristics might increase risk to avian cholera outbreaks due to increased survival of *P. multocida*. The RWB is one of four major U.S. enzootic areas for avian cholera in waterfowl (USGS, 1999a) and previous outbreaks of avian cholera have occurred at McMurtrey NWR and Harvard WPA located within two miles of the created wetlands (Smith et al., 1989). Created wetland water quality characteristics that enhance survival of *P. multocida* in laboratory studies include warmer water temperature, increased protein material (Bredy and Botzler, 1989), and high concentrations of Ca and Mg (Price et al., 1992). In addition, the created wetlands exhibit high conductivity, a condition associated with avian cholera outbreaks in Nebraska wetlands (Windingstad et al., 1984; Gordon, 1989). However, more recent analysis by USGS (1999b) found no associations between risk of avian cholera outbreaks and Ca, Mg, specific conductance, protein, or the abundance of *P. multocida*. Mallard ducks exposed to sewage sludge in their diet did not exhibit increased susceptibility to avian cholera, but exhibited increased cadmium concentrations in liver (Goldberg and Yuill, 1989).

Salmonella appears to be associated with the swine wastewater as it was frequently isolated from samples collected on sites that receive swine wastewater effluent. Salmonellosis has caused die-offs of several captive-reared avian species including waterfowl; although, large-scale die-offs of free ranging wild birds other than songbirds and colonial nesting birds, have rarely been reported (Dr. Kathy Converse, Wildlife Disease Specialist at NWHC, pers. comm., 2004).

Recoveries of *Yersinia intermedia* from all sites indicates that the pathogen is not likely specific to swine wastewater; however, *Yersinia enterocolitica* recoveries only occurred in areas associated with swine waste. The disease potential of *Y. intermedia* in wildlife species is not well understood (Aleksic and Bockemühl, 1999) and yersiniosis in wild waterfowl or shorebirds is not commonly reported to the NWHC (Dr. Kathy Converse, Wildlife Disease Specialist at NWHC, pers. comm., 2004). Potential sources

for the *Yersinia* recoveries at McMurtrey Marsh are warm-blooded mammals, such as cattle that graze the area, as well as non-point source agricultural runoff from Hastings Pork or other swine and cattle feeding operations in McMurtrey NWR's watershed. Further research and pathogen fingerprinting techniques would be needed to determine whether disease pathogens are being transferred to McMurtrey NWR from Hastings Pork and whether the exposure pathways include waterfowl pathogen carriers or sediment and water from run-off.

Bacterial antibiotic resistance was not included in the original study design; however, the finding of multiple antibiotic resistances in *Salmonella* spp. and *E. coli* isolates recovered from the Hastings Pork lagoons, canals, and created wetlands raises a concern. Antibiotic resistance may not present a direct threat to wildlife. Nonetheless, human health concerns related to the spreading of antibiotic resistant bacteria by waterfowl may dictate how these species are managed.

Hormones

Hormone exposure to waterfowl at the created wetlands needs to be further evaluated. The created wetlands appear to be contaminated with testosterone and E₂ from the CAFO waste; however, only a limited number (n=5) of water samples were analyzed and samples were run directly. The limited sampling indicates that the lagoons do not appear to limit concentrations of E₂ and testosterone in the created wetlands and these hormones may lead to adverse effects to wetland wildlife. Concentrations of E₂ in three of four samples from Hastings pork exceeded 10 ng/L, a concentration likely to exert significant adverse effects on wildlife reproduction (Witters et al., 2000). Amphibians may be the species at greatest risk to hormone exposure at the created wetlands. Amphibian exposure to E₂ results in vitellogenin induction in adults (Palmer and Palmer, 1995) and sexual differentiation in larvae (Hayes, 1998). There are no known studies that have evaluated waterfowl or shorebird exposure to natural hormones from ingestion of contaminated water or food items.

Water Quality

Created wetland nutrients, pH, and specific conductivity were significantly greater than those measured on RWB wetlands and, although they did not exceed any known toxicity thresholds for waterfowl, they are resulting in eutrophication. Eutrophication of this system may represent the greatest health risks to waterfowl that utilize the area as previous research indicates wetland eutrophication can adversely affect waterfowl by decreasing their foraging potential (Gaiser and Lang, 1998) or by creating an environment conducive to disease pathogens and toxin-producing algal blooms (USGS, 1999a; Carmichael, 1997). Eutrophication of the created wetlands is likely limiting their potential as quality habitat for waterfowl and shorebirds by altering natural wetland invertebrate and plant communities. Eutrophication in wetland systems is characterized by decreased invertebrate species density and richness, dominance by few nutrient tolerant taxa, and loss of endemic and characteristic species (Bedford et al., 1999). Anoxic sediments from eutrophication are likely the cause for chironomid dominance in the benthic invertebrate communities of the created wetlands and the decreased Shannon-Wiener diversity index. Decreased taxa diversity and dominance of the invertebrate community by chironomids is common in areas of degraded water and sediment quality (Dickman and Rygiel, 1996 and 1998; Chow-Fraser, 1998; Victor and Onomivbori, 1999; Nelson et al., 2000). Wetland eutrophication among prairie pothole wetlands in northwest Iowa also was associated with limited abundance and composition of microcrustaceans (Gaiser and Lang, 1998). Although flying insect abundance was not measured in this study, wetland eutrophication tends to decrease abundance of flying insects when compared to oligotrophic wetlands and may contribute to relatively poor foraging conditions for young waterfowl (King and Brazner, 1999).

Differences in invertebrate communities between the created wetlands and McMurtrey Marsh also may be due to differences in water regimes. McMurtrey Marsh functions as a seasonal wetland, whereas the created wetlands function as open permanent wetlands creating an environment more conducive to chironomid species. In

North Dakota, chironomids were reported to be the dominant fauna of semi-permanent prairie wetlands and comprised a smaller percentage in seasonal wetlands (Nelson, 1990).

Advanced eutrophication can result in bare mud substrates with anoxic sediments, an environment that results in botulism-related mortality of birds (Crowder and Bristow, 1988). Avian botulism also is associated with sewage and other wastewater discharges into marshes (USGS, 1999a). Created wetland water was frequently between a pH of 8 to 10, greater than 20°C, and less than 2 parts per thousand salinity; all conditions that tend to favor avian botulism outbreaks (Rocke and Samuel, 1999).

The environment created by the eutrophication of the created wetlands also is suitable for cyanobacteria blooms and results from this study indicate that waterfowl and shorebirds are likely exposed to cyanobacteria toxins. Cyanobacteria toxins can be lethal to foraging dabbling ducks (Carmichael, 1992; Matsunaga, 1999). In Japan, 20 spot billed ducks (*Anas paecilorhyncha*) died from acute exposure to microcystins in a pond following eutrophication caused by an influx of untreated sewage (Matsunaga, 1999). The Nebraska Game and Parks Commission (NGPC) also have attributed waterfowl die-offs in eastern Nebraska to cyanobacteria blooms (NGPC, 1992). In August and September of 1992, three separate duck die-off events occurred in a small lake at Fontenelle Park in Omaha, Nebraska (NGPC, 1992). After the third die-off occurred, it was confirmed that water samples contained an abundance of *Microcystis* spp. and *Anabaena circinalis*, two species that produce microcystin toxins (NGPC, 1992).

Uncertainty Analysis and Data Gaps

Ongoing research

The effects of CAFO contaminants on waterfowl that utilize the Hastings Pork created wetlands are currently being investigated in a separate study titled “Post-Remediation Evaluation Using Mallard Sentinels at the Hastings Pork Confined Animal Feeding Operation and Implications for Water Quality at McMurtrey NWR.” Waterfowl enclosures were constructed on two of the created wetlands (CW4 and CW6) and on two control RWB wetlands located within three miles of Hastings Pork (McMurtrey NWR

and Harvard WPA). This research will provide further data on water quality parameters for the created wetlands and will measure disease pathogens and concentrations of trace elements and microcystins in mallard sentinels. Concentrations of Se in sentinel mallard eggs and liver, in conjunction with measurements of Se in sediments, water, and invertebrates obtained from this study, will allow for an aquatic hazard assessment of Se as described by Lemly (1995). At the time of this report, field sampling for this project has been completed; however, sample analysis and data interpretation are ongoing.

Future Research Needs

Trace element concentrations in created wetland plants, sediments, water, and invertebrates may continue to accumulate over time as the created wetlands have no outlet and thus serve as a sink for contaminants that enter from the lagoons. In addition, clay soils have a greater capacity to adsorb metals and phosphorus than other soil types and negative effects may develop only after adsorption sites are exhausted after long periods of continued application (Ap Dewi, 1994). Land application of swine waste and subsequent field run-off also can lead to trace element contamination of surface waters (Eisler, 1990). The use of swine waste as a fertilizer can lead to increased inputs of phosphorus in run-off as the soil's capacity to adsorb phosphorus diminishes over time (Racz and Fitzgerald, 2001). Future monitoring will be needed to evaluate these potential delayed effects on trace element accumulation in created wetland biota, sediments, and plants. It is recommended that the Service conduct an aquatic hazard assessment for Se, as described by Lemly (1995), within 5 to 10 years from this report.

Further monitoring is needed to determine if natural hormone concentrations in the created wetlands are high enough to adversely affect wetland species. In addition to E₂ and testosterone, equol (a phytoestrogen) also may be a concern and should be included in any future monitoring of natural hormones at the site. Concentrations of equol as high as 16.6 ppm have been reported in swine manure (Burnison et al., 2002). Although equol is between 200 and 1,000 times less estrogenic than E₂ (Burnison et al., 2002); it may still be an important contributor to the total estrogen exposure.

Future pathogen screening at the site should include *Trichomonas* spp. Swine manure can be a source of trichomoniasis, a disease caused by the protozoan parasite *Trichomonas suis* (Ap Dewi, 1994). Although it is rare in free-ranging waterfowl, it has caused major die-offs in doves and pigeons as well as less visible chronic losses (USGS, 1999a). This may be of importance, as morning doves (*Zenaida macroura*) appear to be attracted to the study area.

The potential for contaminants in swine wastewater to modify RWB wetland invertebrate communities needs to be further researched. Only benthic invertebrates from McMurtrey Marsh, a temporary wetland, were compared to those from the created wetlands. Invertebrate community structures from other RWB wetlands, especially those that with areas that function more as permanent wetlands (e.g., Smith WPA), also should be compared to the created wetlands and RWB wetlands with animal feeding operations in their watershed.

Lagoon water may be an important exposure pathway for CAFO contaminants to waterfowl. Although lagoon water was analyzed for antibiotics and disease pathogens, trace elements concentrations in lagoon water were not determined during this study and should be included in future site evaluations.

Recommendations

In April of 2001, NEFO personnel met with Hastings Pork and the RWBJV to discuss preliminary results. Four management recommendations were agreed upon: 1) water from Hastings Pork would not be used as a supplemental water source for McMurtrey NWR; 2) Phase II of the Hastings Pork/RWBJV partnership, to create additional wetlands in the summer of 2001, would be delayed; 3) water quality would be improved by implementing a comprehensive nutrient management plan that applies “Best Management Practices” including remediation of the effluent delivery system; and 4) a post-remediation evaluation of contaminants at the site and an assessment of contaminant exposure and effects to waterfowl would be produced before construction of additional created wetlands would be supported by the RWBJV.

Remediation work to improve water quality (recommendation # 3 above) scheduled for the summer of 2001 was not completed by Hastings Pork, as they decided to halt any further development of created wetlands or the remediation of existing created wetlands until all current Service research investigations were completed. Consequently, recommended management actions to Hastings Pork will be included in the final report for the mallard enclosure study. These recommendations will likely focus on the use of remediation measures, such as the use of constructed wetlands that are designed specifically to treat domestic sewage to improve the quality of the influent to the created wetland habitat. The treatment of CAFO wastewater is essential in protecting Federal trust resources at McMurtrey NWR as the movement of this effluent into the refuge during heavy rain events remains a concern.

Conclusions

Although lagoon sediments typically contained the highest concentrations of CAFO contaminants and toxicity thresholds for metals were frequently exceeded in lagoons, waterfowl and shorebirds are most likely exposed to CAFO contaminants while foraging in the created wetlands.

Sediment toxicity thresholds for trace elements were exceeded only in lagoons and canals; however, many trace elements that are associated with hog-manure appear to be reaching the created wetlands. Accumulation of these contaminants may increase with time possibly resulting in detrimental effects to wetland plants and invertebrates as well as waterfowl and shorebird species.

Antibiotic exposure to waterfowl and shorebirds in the created wetlands appeared to be essentially negligible, but the number of samples from created wetlands that were analyzed for antibiotics was small and the detection limits may have been too high.

Swine wastewater from Hastings Pork does not appear to be a source for *Erysipelothrix* spp. or *P. multocida*. Bacteria pathogens that appear to be associated with the Hastings Pork swine waste effluent and may be of concern include *Salmonella* spp. and *Yersinia enterocolitica*.

When compared to McMurtrey Marsh and other RWB wetlands, the created wetlands exhibited increased pH, specific conductivity, and salinity. Eutrophication of the created wetlands may represent the greatest health threat to waterfowl that utilize these wetlands by creating an environment that is conducive to cyanobacteria blooms and outbreaks of avian botulism and avian cholera. Large concentrations of waterfowl may be attracted to the created wetlands during drought years when other wetlands in the area are dry, potentially resulting in large-scale waterfowl or shorebird mortality events.

Created wetland invertebrate communities were dominated by abundant populations of pollutant tolerant chironomid species and were less diverse than those in McMurtrey Marsh. Eutrophication also may be degrading wetland habitat quality by limiting plant community diversity.

On-going research will further evaluate CAFO contaminant exposure and effects to waterfowl that utilize the created wetlands. Future research needs are to monitor the accumulation of trace elements in the created wetlands over time to determine if adsorption capacities in sediments and soils that receive swine waste are depleted and to evaluate the adverse effects that natural hormones from swine may have on waterfowl and/or shorebirds that utilize these wetlands.

Recommended remediation strategies will be included in the final report for the follow-up mallard enclosure study, and will likely focus on the use of constructed wetlands to improve water quality before it is used to create waterfowl habitat in the created wetlands. Further treatment is needed to decrease concentrations of contaminants in Hastings Pork swine wastewater before it is used to create waterfowl habitat as “High quality habitat is the key to healthy waterfowl populations” (Friend, 1992).

REFERENCES

- Aleksic S, Bockemühl J. 1999. *Yersinia* and other Enterobacteriaceae. In Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover RH, eds, *Manual of Clinical Microbiology*. ASM Press, Washington, D.C. pp 483-491.
- Ap Dewi I. 1994. The use of animal waste as a crop fertilizer. In Ap Dewi I, Axford RFE, Fayez I, Marai M, Omed H, eds, *Pollution in Livestock Production Systems*. CAB International, UK. pp. 309-332.
- Arzul G, Maguer JF. 1990. Influence of pig farming on the copper content of estuarine sediments in Brittany, France. *Marine Pollution Bulletin* 21:431-434.
- Atienzar FA, Billingham Z, Depledge MH. 4-n-nonylphenol and 17- β estradiol may induce common DNA effects in developing barnacle larvae. *Environmental Pollution* 120:735-738.
- Baker DL, Lusk JD, Giggelman CM. 1998. Phase II contaminants investigation of Buffalo Lake National Wildlife Refuge, Texas 1993-1994. *U.S. Fish and Wildlife Service Region 2 Contaminants Program*. Arlington Ecological Services Field Office. 35 pp.
- Barbour MT, Gerritsen J, Snyder BD, Strubling JB. 1999. Rapid bioassessment protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates and fish, second edition. *EPA 841-B-99-002*. U.S. Environmental Protection Agency; Office of Water; Washington, D.C.
<http://www.epa.gov/owow/monitoring/rbp/>.
- Barnes B, Nudds TD. 1991. Salt tolerance in American black ducks, mallards, and their F1 hybrids. *The Auk* 108:89-98.
- Bedford BL. 1999. Cumulative effects on wetland landscapes: links to wetland restoration in the United States and southern Canada. *Wetlands* 19:775-788.
- Benning DS. 1987. Coordinated mid-continent white-fronted goose survey. *U.S. Fish and Wildlife Service Annual Report*. 7 pp.
- Beyer NW, Day D, Melancon MJ, Sileo L. 2000. Toxicity of Anacostia River, Washington, D.C., USA, sediment fed to mute swans (*Cygnus olor*). *Environmental Toxicology and Chemistry* 19(3):731-735.
- Bortner JB, Johnson FA, Smith GW, Trost RE. 1991. 1991 status of waterfowl and fall flight forecast. *U.S. Fish and Wildlife Service*. 38 pp.
- Bredey JP, Botzler RG. 1989. The effects of six environmental variables on *Pasteurella multocida* population in water. *Journal of Wildlife Disease* 25(2):232-239.
- Buckley JA. 1994. Bioavailability of copper in wastewater to *Lemna minor* with biological and electrochemical measures of complexation. *Water Research* 28(12):2457-2467.
- Burkholder JM, Mallin MA, Glasgow HB Jr., Larsen ML, McIver MR, Shank CG, Deame-Melina N, Briley DS, Springer J, Touchette BW, Hannon EK. 1997. Impacts to a coastal river and estuary from rupture of a large swine holding lagoon. *J Environ Qua* 26:1451-1466.

- Burnison KB, Hartmann A, Lister A, Servos MR, Thomas T, Van Der Kraak G. 2003. A toxicity identification evaluation approach to studying estrogenic substances in hog manure and agricultural runoff. *Environmental Toxicology and Chemistry* 22(10):2243-2250.
- Cain BW, Sileo L, Franson JC, Moore J. 1983. Effects of dietary cadmium on mallard ducklings. *Environ Res* 32:286-297.
- Cain BW, Pafford EA. 1981. Effects of dietary nickel on survival and growth of mallard ducklings. *Archives of Environmental Contamination and Toxicology* 10:737-745.
- Calabrese EJ, Baldwin LA. 1993. *Performing ecological risk assessments*. Lewis Publishers, Ann Arbor, MI. 257 pp.
- Camardese MB, Hoffman DJ, LeCaptain LJ, Pendleton GW. 1990. Effects of arsenate on growth and physiology in mallard ducklings. *Environ Toxicol Chem* 9:785-795.
- Canadian Council of Ministers of the Environment (CCME). 2002. Canadian water quality guidelines for the protection of aquatic life. <http://www.ec.gc.ca/ceqg-rcqe/English/ceqg/water/default.cfm#aqu>.
- Carmichael WW. 1992. A status report on planktonic cyanobacteria (blue-green algae) and their toxins. *Environmental Protection Agency*. EPA/600/R-92/079. 141 pp.
- Carmichael WW. 1997. The cyanotoxins. *Advances in Botanical Research*. 27:211-240.
- Chow-Fraser P. 1998. A conceptual ecological model to aid restoration of Cootes Paradise Marsh, a degraded coastal wetland of Lake Ontario, Canada. *Wetlands Ecology and Management* 6(1):43-57.
- Christie P, Beattie JAM. 1989. Grassland soil microbial biomass and accumulation of potentially toxic metals from long-term slurry application. *Journal of Applied Ecology* 26:597-612.
- Clesceri, LS, Greenberg EA, Eaton AD. 1998. Standard methods for the examination of water and wastewater. *American Public Health Association*, Washington, D.C.
- Crowder AA, Bristow JM. 1988. The future of waterfowl habitats in the Canadian lower Great Lakes wetlands. *Journal of Great Lakes Research* 14(1):115-127.
- Demayo A, Taylor MC, Taylor KW. 1982a. Effects of copper on humans, laboratory and farm animals, terrestrial plants, and aquatic life. *Crit Rev Environ Control* 12:183-255.
- Demayo A, Taylor MC, Taylor KW, Hodson PV. 1982b. Toxic effects of lead and lead compounds on human health, aquatic life, wildlife plants, and livestock. *Crit Rev Environ Control* 12:257-305.
- Denn J. 1999. Animal farms endanger water quality. *Water Technology Online March 1999*. <http://www.watertechonline.com/article.asp?IndexID=5220304>
- Desbrow C, Routledge E, Brighty GC, Sumpter JP, Waldock M. 1998. Identification of estrogenic chemicals in STW effluent: chemical fractionation and in vitro biological screening. *Environmental Science & Technology* 32(11):1549-1558.
- Dickman M, Rygiel G. 1996. Chironomid larval deformity frequencies, mortality, and diversity in heavy-metal contaminated sediments of a Canadian riverine wetland. *Environment International* 22(6):693-703.
- Dickman M, Rygiel G. 1998. Municipal landfill impacts on a natural stream located in an urban wetland in regional Niagara, Ontario. *Canadian Field-Naturalist* 112(4):619-630.

- Ducks Unlimited. 2003. Ducks Unlimited 2003 Annual Report. 25 pp.
<http://www.ducks.org/about/2003AnnualReport/AnnualReport--Main.pdf>
- Dupont HL, Steele JH. 1987. Use of antimicrobial agents in animal feeds: implications for human health. *Reviews of Infectious Diseases* 9(3):447-460.
- Eastin WC Jr., O'Shea TJ. 1981. Effects of dietary nickel on mallards. *Journal of Toxicology and Environmental Health* 7:883-892.
- Echols KR. 2001. Microcystin algal toxins in algal bloom samples from Nebraska wetlands. *U.S. Geological Survey Columbia Environmental Research Center Final Report*, Columbia, MO. 8 pp.
- Eisler R. 1985. Cadmium hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Fish and Wildlife Service Biological Report 85 (1.2), Contaminant Hazard Reviews Report 2*. 46 pp.
- Eisler R. 1988a. Arsenic hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Fish and Wildlife Service Biological Report 85 (1.12), Contaminant Hazard Reviews Report 12*. 92 pp.
- Eisler R. 1988b. Lead hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Fish and Wildlife Service Biological Report 85 (1.14), Contaminant Hazard Reviews Report 14*. 134 pp.
- Eisler R. 1989. Molybdenum hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Fish and Wildlife Service Biological Report 85 (1.19), Contaminant Hazard Reviews Report 19*. 61 pp.
- Eisler R. 1990. Boron hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Fish and Wildlife Service Biological Report 85 (1.20), Contaminant Hazard Reviews Report 20*. 32 pp.
- Eisler R. 1993. Zinc hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Fish and Wildlife Service Biological Report 10, Contaminant Hazard Reviews Report 26*. 106 pp.
- Eisler R. 1998a. Copper hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Geological Survey Biological Science Report USGS/BRD/BSR--1997-0002, Contaminant Hazard Reviews Report 33*. 98 pp.
- Eisler R. 1998b. Nickel hazards to fish, wildlife, and invertebrates: a synoptic review. *U.S. Geological Survey Biological Science Report USGS/BRD/BSR--1998-0001, Contaminant Hazard Reviews Report 34*. 76 pp.
- Eldridge J. 1990. Aquatic invertebrates important for waterfowl production. In Cross DH (ed) *Waterfowl Management Handbook*. U. S. Fish and Wildlife Service, Washington, DC, USA.
<http://www.nwrc.usgs.gov/wdb/pub/wmh/contents.html>
- European Agency for the Evaluation of Medicinal Products (EAEM). 1995. Committee for veterinary medicinal products oxytetracycline, tetracycline, chlortetracycline. *Maximum Residue Limits Summary Reports 1-3*.
<http://www.emea.eu.int/htms/vet/mrls/m-rmrl.htm>.
- European Commission. 1999. Assessment of potential risks to human health from hormone residues in bovine meat and meat products. XXIV/B3/SC4. 136 pp. http://europa.eu.int/comm/food/fs/sc/scv/out21_en.pdf.
- Falconer IR, Smith JV, Jackson ARB, Jones A, Runnegar MT. 1988. Oral toxicity of a bloom of the cyanobacterium *Microcystis aeruginosa* administered to mice over periods up to one year. *J Toxicol Environ Health* 24:291-305.
- Falconer IR. 1991. Tumor promotion and liver injury caused by oral consumption of cyanobacteria. *Environmental Toxicology and Water Quality: An International Journal* 6:177-184.

- Finlay-Moore O, Hartel PG, Cabrera ML. 2000. 17 beta-estradiol and testosterone in soil and runoff from grasslands amended with broiler litter. *Journal of Environmental Quality* 29(5):1604-1611.
- Fraser H. 1991. Manure characteristics fact sheet. *Ontario Ministry of Agriculture, Food, and Rural Affairs. Agdex # 538*, www.gov.on.ca/OMAFRA/english/livestock/swine/facts/85-109.htm.
- Friend M. 1992. Environmental influences on major waterfowl diseases. *Trans 57th N.A. Wild & Nat Res Conf* 517-52.
- Friend M. 2002. Avian disease at the Salton Sea. *Hydrobiologia* 473(1-3):293-306.
- Furness RW. 1996. Cadmium in birds. In Beyer WN, Heinz GH, and Redmon-Norwood AW, eds *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*. CRC Press, Lewis Publishers, Boca Raton, Florida. pp 389-405.
- Gaiser EE, Lang KL. 1998. Distribution of cladoceran zooplankton among prairie pothole wetlands in northwest Iowa. *Lake and Reservoir Management* 14(1):37-51.
- Gensemer RW, Playle RC. 1999. The bioavailability and toxicity of aluminum in aquatic environments. *Critical Reviews in Environmental Science and Technology* 29(4):315-450.
- Gersib RA, Dinan KF, Kauffeld JD, Onnen MD, Gabig PJ, Cornely JE, Jasmer GE, Hyland JM, Strom KJ. 1992. *Rainwater Basin Joint Venture Implementation Plan*. Nebraska Game and Parks Commission, Lincoln, NE. 56 pp.
- Gersich, FM. 1984. Evaluation of a static renewal chronic toxicity test method for *Daphnia magna* Straus using boric acid. *Environmental Toxicology and Chemistry* 3(1):89-94.
- Goldberg DR, Yuill TM. 1989. Effects of sewage sludge on the immune defenses of mallards. *Environmental Research* 51:209-217.
- Goñi-Urriza M, Capdepuuy M, Arpin C, Raymond N, Caumette P, Quentin C. Impact of an urban effluent on antibiotic resistance of riverine Enterobacteriaceae and *Aeromonas* spp. *Applied and Environmental Microbiology* 66(1):125-132.
- Gordon CC. 1989. The relationship of wetland characteristics to avian cholera (*Pasteurella multocida*) outbreaks in the Rainwater Basin area of Nebraska. *Master's thesis submitted to South Dakota State University*. 126 pp.
- Goyer RA, Clarkson TW. 2001. Toxic effects of metals. In Klaassen CD, ed, *6th Edition Casarett & Doull's Toxicology: The Basic Science of Poisons*. McGraw-Hill Medical publishing division. p. 811-867.
- Gross TS, Arnold BS, Sepúlveda, McDonald K. 2003. Endocrine disrupting chemicals and endocrine active agents. In Hoffman DJ, Rattner BA, Burton GA Jr., Cairns J Jr., eds, *Handbook of Ecotoxicology Second Edition*. CRC Press. Pages 1033-1098.
- Guardabassi L, Petersen A, Olsen JE, Dalsgaard A. Antibiotic resistance in *Acinetobacter* spp. isolated from sewers receiving waste effluent from a hospital and a pharmaceutical plant. *Applied and Environmental Microbiology* 64(9):3499-3502.
- Gupta UC, Jame YW, Campbell CA, Leyshon AJ, Nicholaichuk W. 1985. Boron toxicity and deficiency: a review. *Can J Soil Sci* 65:381-409.

- Halling-Sørensen, B, Nors Nielsen S, Lanzky PF, Ingerslev F, Holten Lützhøft HC, Jørgensen SE. 1997. Occurrence, Fate and Effects of Pharmaceutical substances in the environment: a review. *Chemosphere* 36(2):357-393.
- Harries JE, Janbakhsh A, Jobling S, Matthiessen P, Sumpter JP, Tyler CR. 1999. Estrogenic potency of effluent from two sewage treatment works in the United Kingdom. *Environmental Toxicology and Chemistry* 18(5):932-937.
- Hatakeyama S. 1999. Effect of copper and zinc on the growth and emergence of *Epeorus latifolium* (Ephemeroptera) in an indoor model stream. *Hydrobiologia* 174:17-27.
- Hayes TB. 1998. Sex determination and primary sex differentiation in amphibians: genetic and developmental mechanisms. *J Exp Zool* 281:373-399.
- Hazardous Substances Data Bank (HSDB). National Library of Medicine, Bethesda, MD (CD-ROM Version), *Micromedex, Inc, Englewood, CO, 2001* <http://toxnet.nlm.nih.gov/>.
- Heinz GH. 1996. Selenium in birds. In Beyer WN, Heinz GH, Redmon AW, eds, *Interpreting Environmental Contaminants in Animal Tissues*. Lewis Publishers, Boca Raton, Florida. pp 453-464.
- Hull RN, Suter GW II. 1994. Toxicological benchmarks for screening contaminants of potential concern for effects on sediment-associated biota: 1994 revision. *Rept. ES/ER/TM-95-R1*. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Irwin LK, Gray S, Oberdorster E. 2001. Vitellogenin induction in painted turtle *Chrysemys picta*, as a biomarker of exposure to environmental levels of estradiol. *Aquat Toxicol* 55:49-60.
- Jensen WI, Cotter SE. 1976. An outbreak of erysipelas in eared grebes (*Podiceps nigricollis*). *Journal of Wildlife Diseases* 12:583-586.
- Jones DS, Suter GW II, Hull RN. 1997. Toxicological benchmarks for screening potential contaminants of concern for effects on sediment-associated biota: 1997 revision. *ES/ER/TM-95/R3*. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Kelly CP, Pothoulakis C, LaMont JT. 1994. *Clostridium difficile* colitis. *New England Journal of Medicine* 330(4):257-262.
- Kelly ME, Fitzgerald SD, Aulerich RJ, Balander RJ, Powell DC, Stickle RL, Stevens W, Cray C, Tempelmn RJ, Bursian SJ. 1998. Acute effects of lead, steel, tungsten-iron, and tungsten-polymer shot administered to game-farm mallards. *Journal of Wildlife Diseases* 43(4):673-687.
- King RS, Brazner JC. 1999. Coastal wetland insect communities along a trophic gradient in Green Bay, Lake Michigan. *Wetlands* 19(2):426-437.
- Koller LD, Exon JH. 1986. The two faces of selenium-deficiency and toxicity are similar in animals and man. *Can J Vet Res.*50:297-306.
- Kolpin DW, Furlong ET, Meyer MT, Thurman ME, Zaugg SD, Barber LB, Buxton HT. Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. Streams, 1999-2000: A national Reconnaissance. *Environmental Science and Technology* 36:1202-1211.
- Krantzberg G. 1989. Metal accumulation by chironomid larvae: the effects of age and body weight on metal body burdens. *Hydrobiologia* 188/189:497-506.

- Lange IG, Daxenberger A, Schiffer B, Witters H, Ibarreta D, Meyer HHD. 2002. Sex hormones originating from different livestock production systems: fate and potential disrupting activity in the environment. *Analytica Chimica Acta* 473:27-37.
- Lee HB, Liu D. 2002. Degradation of 17 beta-estradiol and its metabolites by sewage bacteria. *Water, Air, & Soil Pollution* 134(1-4):351-366.
- Lemly DA. 1995. A protocol for aquatic hazard assessment of selenium. *Ecotoxicology and Environmental Safety* 32:280-288.
- Ley DH. 1986. Nitrite poisoning in herring gulls (*Larus argentatus*) and ring-billed gulls (*Larus delawarensis*). *Journal of Wildlife Diseases* 22(3):381-384.
- Long ER, MacDonald DD, Smith SL, Calder FD. 1995. Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environ. Manage.* 19:81-97.
- MacDonald DD, Ingersoll CG, Berger TA. 2000. Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Archives of Environmental Contamination and Toxicology* 39:20-31.
- Mallin MA, Burkholder JM, McIver, MR, Shank CG, Glasgow HB Jr., Touchette BW, Springer J. Comparative effects of poultry and swine waste lagoon spills on the quality of receiving stream waters. *J. Environmental Quality* 26:1622-1631.
- Marco A, Blaustein AR. 1999. The effects of nitrite on behavior and metamorphosis in cascades frogs (*Rana cascadae*). *Environmental Toxicology and Chemistry* 18(5):946-949.
- Marco A, Quilchano C, Blaustein AR. 1999. Sensitivity to nitrate and nitrite in pond-breeding amphibians from the Pacific Northwest, USA. *Environmental Toxicology and Chemistry* 18:(12)2836-2839.
- Matsunaga H, Harada KI, Senma M, Ito Y, Yasuda N, Ushida S, Kimura Y. 1999. Possible cause of unnatural mass death of wild birds in a pond in Nishinomiya, Japan: sudden appearance of toxic cyanobacteria. *Natural Toxins*. 7(2):81-84.
- Mitcham SA, Wobeser, G. 1988. Effects of sodium and magnesium sulfate in drinking water on mallard ducklings. *Journal of Wildlife Diseases* 24:30-44.
- Murphy TP, Lawson A, Nalewajko C, Murkin H, Ross L, Oguma K, McIntyre T. 2000. Algal toxins-initiators of avian botulism? *Environ Toxicol* 15:558-567.
- National Academy of Sciences (NAS). 1975. Medical and biological effects of environmental pollutants: nickel. *National Research Council, National Academy of Sciences, Washington, D.C.* 277 pp.
- National Agricultural Statistics Service (NASS). 1996. *Nebraska agricultural statistics 1995-1996*. Issued October 1996. http://www.nass.usda.gov/ne/9596book/pag_090.htm
- National Agricultural Statistics Service (NASS). 2002. *Nebraska agri-facts special edition*. Issued April 2002. <http://www.nass.usda.gov/ne/special/agfspc02.pdf>
- National Irrigation Water Quality Program (NIWQP). 2003. *Data Synthesis. Sun River Area, Montana*. http://www.usbr.gov/niwqp/data/study_areas/sun.htm
- Nebraska Department of Environmental Quality (NDEQ). 1997. *Wetland water quality monitoring-1997*. 23 pp.

- Nebraska Department of Environmental Quality (NDEQ). 2002. *Title 117 Nebraska Water Quality Standards*. <http://www.deq.state.ne.us/>
- Nebraska Department of Environmental Quality (NDEQ). 2003. Fish kills reported by NDEQ 1996-2003. Unpublished data.
- Nebraska Game and Parks Commission (NGPC). 1992. Wildlife disease and mortality investigations. *Pittman-Robertson Project W-15-R-49*. 19 pp.
- Nelson RD. 1990. Seasonal abundance and life cycles of chironomids (Diptera:Chironomidea) in four prairie wetlands. *Dissertation Abstracts International Part B: Science and Engineering* 51(1):1-124.
- Nelson SM, Roline RA, Thullen JS, Sartoris JJ, Boutwell JE. 2000. Invertebrate assemblages and trace element bioaccumulation associated with constructed wetlands. *Wetlands* 20(2):406-415.
- North American Waterfowl Management Plan. 2003. Strengthening the biological foundations: 2003 strategic guidance. 22 pp. <http://birdhabitat.fws.gov/NAWMP/images/Strategicguidances.pdf>
- Pais I, Jones JB. 1997. *The Handbook of Trace Elements*. St. Lucie Press, Boca Raton, F., 223 pp.
- Palawski DU, Malloy JC, DuBois KL. 1991. Montana national wildlife refuges: contaminant issues of concern. *U.S. Fish and Wildlife Service Progress Report*. Montana State Office, Helena, MT. 96 pp.
- Palmer BD, Palmer SK. 1995. Vitellogenin induction by xenobiotic estrogens in the red-eared turtle and African clawed frog. *Environmental Health Perspectives* 103:19-25.
- Payne GG, Martens DC, Kornegay ET, Lindermann MD. 1988. Availability and form of copper in three soils following eight annual applications of copper-enriched swine manure. *J Environ Qual* 17(4):740-746.
- Popoff MY, Minor LL. 1997. Antigenic formulas of the *Salmonella* serovars. *WHO Collaborating Centre for Reference and Research on Salmonella*, Paris, France.
- Price JI, Brand CJ. 1984. Persistence of *Pasteurella multocida* in Nebraska wetlands under epizootic conditions. *Journal of Wildlife Diseases* 20:90-94.
- Price JI, Yandell BS, Porter WP. 1992. Chemical ions affect survival of avian cholera organisms in pondwater. *Journal of Wildlife Management* 56(2):274-278.
- Quinn PJ, Carter ME, Markey B, Carter GR. 1994. *Clinical Veterinary Microbiology*. Wolfe Publishing, London, pp. 916.
- Racz GJ, Fitzgerald MM. 2001. Nutrient and heavy metal contents of hog manure– effect on soil quality and productivity. *Livestock Options for the Future. National Conference* in Winnipeg, Manitoba, Canada, June 25-27. 21 pp. <http://www.gov.mb.ca/agriculture/livestock/livestockopt/papers/racz.pdf>
- Rainwater Basing Joint Venture (RWBJV). 2000. The Rainwater Basin Joint Venture evaluation plan July 2000. 55 pp. http://www.rwbjv.org/pdf/RWB_JV_Eval_Plan.pdf

- Rocke TE, Samuel MD. 1999. Water and sediment characteristics associated with avian botulism outbreaks in wetlands. *Journal of Wildlife Management* 63(4):1249-1260.
- Rodriguez RE, Misra M, Diwan BA, Riggs CW, Kasprzak KS. 1996. Relative susceptibilities of C57BL/6X, (C57BL/6 X C3H/He)F1, and C3H/He mice to acute toxicity and carcinogenicity of nickel subsulfide. *Toxicology* 107:131-140.
- Rodgers-Gray TP, Jobling S, Morris S, Kelly C, Kirby S, Janbakhsh A, Harries JE, Waldock MJ, Sumpter JP, Tyler CR. 2000. Long-term temporal changes in the estrogenic composition of treated sewage effluent and its biological effects on fish. *Environmental Science & Technology* 34(8):1521-1528.
- Rosen MN, Bischoff AI. 1950. The epidemiology of fowl cholera as it occurs in the wild. *Transactions of the 15th North American Wildlife Conference* 147-154.
- Routledge EJ, Sheahan D, Desbrow C, Brighty GC, Waldock M, Sumpter JP. 1998. Identification of estrogenic chemicals in STW effluent: in vivo responses in trout and roach. *Environmental Science & Technology* 32(11):1559-1565.
- Sanders JG, Cibik SJ. 1985. Adaptive behavior of euryhaline phytoplankton communities to arsenic stress. *Mar Ecol Prog Ser* 22:199-205.
- Schiewer S, Meuser K, Wintgens T. 2001. Process aspects of the behavior of endocrine disrupters in sewage treatment plants. *Wasser und Boden* 53(1-2):10-15.
- Shacklette HT, Boerngen JG. 1984. Element concentrations in soils and other surficial materials of the conterminous United States. *Professional Paper 1270. U.S. Geological Survey, Washington D.C.*
- Sharpley AN, Daniel T, Sims T, Lemunyon J, Stevens R, Parry R. 1999. Agricultural phosphorus and eutrophication. *USDA-Agricultural Research Service, ARS-149, Washington, DC.*
- Shore LS, Correl DL, Chakraborty PK. 1995. Relationship of fertilization with chicken manure and concentration of estrogens in small streams. In Steele K, ed, *Animal Waste and the Land Water Interface*. CRC, Boca Raton, FL, USA, pp 155-162.
- Silva J Jr., Fekety R. 1981. Clostridia and antimicrobial enterocolitis. *Annual Review of Medicine* 32:327-333.
- Sims TJ. 1995. Characteristics of animal wastes and waste-amended soils: an overview of the agricultural and environmental issues. In Steel K, ed, *Animal Waste and the Land-water Interface*. Lewis Publishers, Boca Raton, FL.
- Sivonen, K. 1996. Cyanobacterial toxins and toxin production. *Phycologia* 36(6):12-24.
- Skorupa JP. 1998. Risk assessment for the biota database of the National Irrigation Water Quality Program. *Report to the National Irrigation Water Quality Program (NIWQP)*. U.S. Fish and Wildlife Service, Sacramento, California.
- Snow DD, Cassada DA, Monson SJ, Zhu J, Spalding RF. 2003. Tetracycline and macrolide antibiotics: trace analysis in water and wastewater using solid phase extraction and liquid chromatography-tandem mass spectrometry. pp 161 – 174. In Thurman EM, Ferrer IM, eds, *Using Liquid Chromatography/Mass Spectrometry, (MS/MS), and Time of Flight MS in Analysis of Emerging Contaminants*. Symposium Series 850, American Chemical Society, Washington, D.C.

- Smith BJ, Higgins KF, Gritzner CF. 1989. Land use relationships to avian cholera outbreaks in the Nebraska Rainwater Basin Area. *Prairie Nat* 21(3):125-136.
- Smith GR, Oliphant JC, White WR. 1982. *Clostridium botulinum* type C on the Mersey Estuary. *Journal of Hygiene* 89:507-511.
- Smith L, Sugiyama H. 1988. *Botulism: The Organism, its Toxins, the Disease*. Charles C. Thomas, Springfield, Illinois, USA.
- Sole M, De Alda MJL, Castillo M, Porte C, Ladegaard-Pedersen K, Barcelo D. 2000. Estrogenicity determination in sewage treatment plants and surface waters from the Catalonian Area (NE Spain). *Environmental Science & Technology* 34(24):5076-5083.
- Soto AM, Calabro JM, Precht NV, Yau AY, Orlando EF, Daxenberger A. 2004. Androgenic and estrogenic activity in water bodies receiving cattle feedlot effluent in eastern Nebraska, USA. *Environ Health Perspect* 112:346-352.
- Sparling, DW, Lowe PT, Campbell PGC. 1997. In Yokel RA, Golub MS, eds, *Ecotoxicology of Aluminum in Research issues in Aluminum Toxicity*. Taylor and Frances Publishers. Washington D.C. pp 47-68.
- Stanley TR Jr., Spann JW, Smith GJ, Rosscoe R. 1994. Main and interactive effects of arsenic and selenium on mallard reproduction and duckling growth and survival. *Arch Environ Contam Toxicol* 26:444-451.
- Statistical Analysis System (SAS) Institute. 2001. *Proprietary Software Release 8.2*: Cary, NC, USA.
- Stolley DS, Bissonette JA, Kadlec JA. 1999. Effects of saline environments on the survival of wild goslings (*Branta canadensis*). *American Midland Naturalist* 142(1):181-190.
- Stubbs K, Cathey K. 1999. CAFOs feed a growing problem. *Endangered Species Bulletin* 24(1):14-15.
- Suter GW II, Tsao CL. 1996. Toxicological benchmarks for screening potential contaminants of concern for effects on aquatic biota: 1996 revision. *Risk Assessment Program Health Sciences Research Division Oak Ridge, Tennessee*. 151 pp.
- Thurman EM, Dietze JE, Scribner EA. 2002. Occurrence of antibiotics in water from fish hatcheries. *USGS Fact Sheet* 120-02. 4pp.
- Tolls J. 2001. Sorption of veterinary pharmaceuticals in soils: a review. *Environmental Science and Technology* 35(17):3397-3406.
- Tuttle PL, Janik CA, Wiemyer SN. 1996. Stillwater National Wildlife Refuge wetlands contaminant monitoring. *Region 1 Study ID:93-1-ICO2A*. U.S. Fish and Wildlife Service. 67 pp + Apped.
- U.S. Department of the Interior (USDI). 1998. Guidelines for interpretation of the biological effects of selected constituents in biota, water and sediment. *National irrigation water quality program information report No. 3*. 198 pp. + App. <http://www.usbr.gov/nwqpp>
- U. S. Environmental Protection Agency (USEPA). 1975. Preliminary investigation of effects on the environment of boron, indium, nickel, selenium, tin, vanadium and their compounds. Vol. 1. Boron. *U.S. Environ. Prot. Agency Rep. 56/2-75-005A*. 111 pp.
- U. S. Environmental Protection Agency (USEPA). 1980. Ambient water quality criteria for nickel. *EPA Report 440/5-80-060*. 206 pp.

- U. S. Environmental Protection Agency (USEPA). 1998. Environmental impacts of animal feeding operations. *Preliminary Data Summary: Feedlots Point Source Category Study December 31, 1998*. Office of Water. <http://www.epa.gov/ostwater/guide/feedlots/envimpct.pdf>
- U. S. Environmental Protection Agency (USEPA). 2001. Environmental assessment of proposed revisions to the national pollutant discharge elimination system regulation and the effluent guidelines for concentrated animal feeding operations. *EPA-821-B-01-001*. <http://www.epa.gov/ost/guide/cafo/envir.html>
- U. S. Environmental Protection Agency (USEPA). 2003. National pollutant discharge elimination system permit regulation and effluent limitation guidelines and standards for concentrated animal feeding operation (CAFOs). *Federal Register* 68(29):7177-7269. <http://cfpub.epa.gov/npdes/afo/cafofinalrule.cfm>
- U.S. Department of Agriculture (USDA), U.S. Environmental Protection Agency (USEPA). 2003. Statement of involvement: U.S. Environmental Protection Agency (EPA) and U.S. Department of Agriculture (USDA) cooperation on the implementation of the Clean Water Act regulations for concentrated animal feeding operations. http://www.epa.gov/npdes/pubs/cafo_statement_involvement.pdf
- U.S. Fish and Wildlife Service. 2001. Rainwater Basin Wetland Management District. *Waterfowl Production Areas* http://mountain-prairie.fws.gov/rainwater/waterfowl_production_areas.htm
- U. S. Geological Survey (USGS). 1999a. Field manual of wildlife diseases: general field procedures and diseases in birds. Friend M, Franson JC, eds, *Information and Technology Report 1999-001*. 426 pp. http://www.nwhc.usgs.gov/pub_metadata/field_manual/field_manual.html.
- U. S. Geological Survey (USGS). 1999b. Physical and biological factors associated with avian cholera in wetland ecosystems. *National Wildlife Health Center Research Project Summary Report (July, 1999)*. 15 pp.
- U. S. Geological Survey (USGS). 2001. Investigation of potential bacterial pathogens and their impacts associated with confined animal feeding operations (CAFO) and their potential impacts to a national wildlife refuge in Nebraska. *Draft Completion Report by the National Wildlife Health Center, Madison, WI*. 29pp.
- Victor R, Onomivbori O. 1999. The effects of urban perturbations on the macrobenthic invertebrates of a southern Nigerian stream. *Tropical Limnology* 3:187-204
- Vighi M. 1981. Lead uptake and release in an experimental trophic chain. *Ecotoxicol Environ Safety* 5:177-193.
- Vuori KM. 1995. Direct and indirect effects of iron on river ecosystems. *Annals Zoologici Fennici* 32(3):317-329.
- Wellnitz TA, Grief KA, Sheldon SP. 1994. Response of macroinvertebrates to blooms of iron-depositing bacteria. *Hydrobiologia* 281(1):1-17.
- White DH, Finley MT, Ferrell JF. 1978. Histopathologic effects of dietary cadmium on kidneys and testes of mallard ducks. *J Toxicol Environ Health* 4:551-558.
- Wilber CG. 1980. Toxicology of selenium: A review *Clin Toxicol* 17:171-230.
- Williams T. 1998. Assembly line swine. *Audubon Magazine* March/April p. 26.
- Windingstad RM, Hurt JJ, Trout AK, Cary J. 1984. Avian cholera in Nebraska's rainwater basin. *Trans N Am Wildl Conf* 49:576-583.

- Witters HE, Vangenechten C, Berckmans P. 2000. Detection of estrogenic activity in Flemish surface waters using an in vitro recombinant assay with yeast cells. *Water Science & Technology* 43(2):117-123.
- Wollenberger L, Halling-Sørensen B, Kusk KO. 2000. Acute and chronic toxicity of veterinary antibiotics to *Daphnia magna*. *Chemosphere* 40:723-730.
- Ying G-G, Kookana RS, RuY-J. 2002. Occurrence and fate of hormone steroids in the environment. *Environment International* 28(6):545-551.
- Zhu J, Snow DD, Cassada DA, Monson SJ, Spalding RF. 2001. Analysis of oxytetracycline, tetracycline, and chlortetracycline in water using solid-phase extraction and liquid chromatography tandem mass spectrometry. *Journal of Chromatography A* 928:177-186.
- Zinkl JG, Dey N, Hyland JM, Hurt JJ, Heddleston KL. 1977. An epornitic of avian cholera in waterfowl and common crows in Phelps County, Nebraska, in the spring, 1975. *Journal of Wildlife Diseases* 13(2):194-198.

APPENDIX: ADDITIONAL TABLES

Table A.1. Summary statistics for concentrations of trace elements in water samples from Rainwater Basin wetlands and Hastings Pork created wetlands and canals, Clay County, Nebraska.

| Trace Element | MDL | Canals | | | Created Wetlands | | | Rainwater Basin Wetlands | | |
|---------------|---------|--------------------------------|----------------|---------------|--------------------------------|----------------|----------------|--------------------------------|-----------------|-----------------|
| | | N _D /N _A | Mean ± S.E. | Range | N _D /N _A | Mean ± S.E. | Range | N _D /N _A | Mean ± S.E. | Range |
| Al | <0.1 | 3/3 | 0.4513 ± 0.219 | 0.154 - 0.879 | 5/6 | 5.9666 ± 2.588 | 0.173 - 14.7 | 11/11 | 2.1073 ± 1.2051 | 0.091 - 11.9 |
| As | <0.0056 | 2/3 | 0.0063 ± 0.001 | 0.005 - 0.008 | 6/6 | 0.0167 ± 0.003 | 0.0082 - 0.029 | 9/18 | 0.0035 ± 0.0013 | 0.0006 - 0.0119 |
| B | <0.111 | 3/3 | 0.443 ± 0.054 | 0.373 - 0.549 | 6/6 | 0.4182 ± 0.077 | 0.204 - 0.707 | 10/11 | 0.0425 ± 0.0057 | 0.005 - 0.0718 |
| Ba | <0.001 | 3/3 | 0.039 ± 0.006 | 0.033 - 0.051 | 6/6 | 0.1601 ± 0.048 | 0.0438 - 0.335 | 11/11 | 0.1952 ± 0.0443 | 0.0478 - 0.59 |
| Be | <0.0011 | 0/3 | NA | NA | 1/6 | 0.000691 | NA | 4/11 | 0.0022 ± 0.0004 | 0.0012 - 0.0032 |
| Cd | <0.0011 | 1/3 | 0.0013 | NA | 1/6 | 0.0014 | NA | 4/11 | 0.002 ± 0.0006 | 0.0009 - 0.0036 |
| Cr | <0.0056 | 1/3 | 0.00266 | NA | 3/6 | 0.0086 ± 0.003 | 0.0034 - 0.014 | 4/11 | 0.0082 ± 0.0013 | 0.005 - 0.0113 |
| Cu | <0.005 | 3/3 | 0.054 ± 0.026 | 0.023 - 0.106 | 6/6 | 0.0218 ± 0.002 | 0.016 - 0.029 | 5/11 | 0.0216 ± 0.01 | 0.0062 - 0.0572 |
| Fe | <0.02 | 3/3 | 0.7457 ± 0.266 | 0.261 - 1.18 | 6/6 | 4.0815 ± 2.076 | 0.171 - 13.7 | 10/11 | 2.1403 ± 1.2434 | 0.0457 - 12.2 |
| Hg | <0.005 | 0/3 | NA | NA | 0/6 | NA | NA | 0/18 | NA | NA |
| Mg | <0.02 | 3/3 | 44.6 ± 6.502 | 31.6 - 51.4 | 6/6 | 41.767 ± 5.225 | 25.2 - 55.8 | 11/11 | 9.02 ± 1.4111 | 2.01 - 18.9 |
| Mn | <0.0056 | 3/3 | 0.1707 ± 0.068 | 0.095 - 0.307 | 6/6 | 0.3629 ± 0.122 | 0.0856 - 0.874 | 9/11 | 0.121 ± 0.0918 | 0.002 - 0.85 |
| Mo | <0.002 | 2/3 | 0.011 ± 0.006 | 0.005 - 0.017 | 2/6 | 0.0109 ± 0.004 | 0.0073 - 0.014 | 3/11 | 0.0078 ± 0.0009 | 0.0062 - 0.0094 |
| Ni | <0.0056 | 3/3 | 0.0378 ± 0.003 | 0.034 - 0.044 | 6/6 | 0.0394 ± 0.008 | 0.0233 - 0.074 | 4/11 | 0.0138 ± 0.0028 | 0.0056 - 0.0172 |
| Pb | <0.01 | 2/3 | 0.0048 ± 0.004 | 0.001 - 0.009 | 4/6 | 0.0115 ± 0.005 | 0.003 - 0.025 | 5/11 | 0.0167 ± 0.0074 | 0.0053 - 0.0448 |
| Se | <0.0056 | 1/3 | 0.00351 | NA | 2/6 | 0.0048 ± 0.002 | 0.0027 - 0.007 | 14/18 | 0.006 ± 0.0014 | 0.0013 - 0.0199 |
| Sr | <0.002 | 3/3 | 0.3077 ± 0.068 | 0.22 - 0.442 | 6/6 | 0.2495 ± 0.027 | 0.182 - 0.366 | 11/11 | 0.2572 ± 0.0441 | 0.042 - 0.542 |
| V | <0.0056 | 3/3 | 0.0095 ± 0.004 | 0.006 - 0.017 | 6/6 | 0.0236 ± 0.006 | 0.0086 - 0.046 | 8/11 | 0.0133 ± 0.0026 | 0.0054 - 0.026 |
| Zn | <0.0111 | 3/3 | 0.2567 ± 0.155 | 0.101 - 0.566 | 6/6 | 0.0565 ± 0.011 | 0.013 - 0.094 | 3/11 | 0.0471 ± 0.0205 | 0.0258 - 0.0881 |

Note: MDL = the minimum detection limit; N_D/N_A = the number of samples with detected concentrations over the number of samples analyzed; and NA = not applicable. A single value in the “Mean ± S.E.” column indicates the concentration detected in one sample. The summary statistics are for samples that were above detection limits only.

Table A.2. Monthly counts of fecal coliforms and fecal streptococci colony forming units per 100 ml of water from sites at Hastings Pork and McMurtrey Marsh, Clay County, Nebraska, 2000.

| Pathogen | Site | Monthly count per 100 ml | | |
|-----------------|--------------------|--------------------------|---------|---------|
| | | April | June | October |
| Fecal Coliforms | Lagoon 1 | TNTC | 209,500 | 770,000 |
| | Lagoon 2 | 10,550 | 10,000 | 11,000 |
| | Lagoon 3 | 160,000 | 4,000 | 59,000 |
| | Lagoon 4 | TNTC | 250,000 | TNTC |
| | Canal 1 | 2450 | 1022 | 3866 |
| | Canal 2 | 400 | 2181 | 100 |
| | Canal 3 | 3900 | 9733 | 90,000 |
| | Canal 4 | 4250 | 933 | 2900 |
| | CW 1 | 100 | 1766 | 1875 |
| | CW 2 | 200 | 3340 | 380 |
| | Ditch | ND | 79 | 415 |
| | MM 1 | 50 | ND | 2200 |
| | MM 2 | 150 | ND | 2050 |
| | MM 3 | 150 | ND | 2650 |
| | Fecal Streptococci | Lagoon 1 | 50,000 | 90,500 |
| Lagoon 2 | | 9200 | 5000 | TNTC |
| Lagoon 3 | | 9350 | 3250 | TNTC |
| Lagoon 4 | | 92,000 | 143,500 | 6000 |
| Canal 1 | | 1800 | 4800 | 4250 |
| Canal 2 | | 1700 | 12,100 | 600 |
| Canal 3 | | 9500 | TNTC | 5800 |
| Canal 4 | | 6150 | 1650 | 4100 |
| Wetland 1 | | 200 | TNTC | 2270 |
| Wetland 2 | | 9000 | 125,000 | 890 |
| Ditch | | ND | 258 | 35 |
| MM 1 | | 27 | ND | 300 |
| MM 2 | | 5 | ND | 50 |
| MM 3 | | 8 | ND | 100 |

Note: CW = created wetland, MM = McMurtrey Marsh, Ditch = ditch that drains runoff from Hastings Pork into McMurtrey Marsh, ND = work not done due to dry conditions, TNTC = too numerous to count (i.e., the agar plates were too overgrown to distinguish separate colony units). See Figure 4 for the locations of these sites.

Table A.3. *Salmonella* species (serotype) recovered from Hastings Pork lagoons, created wetlands and canals, Clay County, Nebraska, 2000.

| Site | Date | Source | <i>Salmonella</i> species (serotype) identification |
|----------|--------------|--------|---|
| Lagoon 2 | October 2000 | water | Newport |
| Lagoon 2 | October 2000 | water | Newport |
| Lagoon 2 | October 2000 | water | Newport |
| Lagoon 3 | October 2000 | water | Newport |
| Lagoon 3 | October 2000 | water | Typhimurium (Copenhagen) |
| Lagoon 3 | October 2000 | water | Typhimurium (Copenhagen) |
| Canal 3 | October 2000 | water | Derby |
| Canal 3 | April 2000 | water | Infantis |
| Canal 3 | October 2000 | water | Infantis |
| Canal 1 | October 2000 | water | Newport |
| Canal 1 | October 2000 | water | Newport |
| CW 5 | June 2000 | water | Muenchen |
| CW 1 | October 2000 | water | Newport |
| CW 1 | October 2000 | water | Newport |
| CW 2 | October 2000 | water | Newport |
| Ditch | October 2000 | water | Muenchen |

Note: CW = Hastings Pork created wetland, The “Ditch” drains runoff from Hastings Pork into MM Marsh.

Table A.4. *Yersinia* species recovered from McMurtrey Marsh and Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000.

| Site | Date | Source | <i>Yersinia</i> species identification |
|----------|--------------|----------|---|
| Lagoon 3 | April 2000 | sediment | <i>Yersinia enterocolitica</i> |
| Lagoon 1 | October 2000 | sediment | <i>Yersinia</i> sp. |
| Lagoon 4 | April 2000 | water | <i>Yersinia</i> sp. |
| Canal 2 | April 2000 | sediment | <i>Yersinia intermedia</i> |
| Canal 2 | April 2000 | sediment | <i>Yersinia intermedia</i> |
| Canal 1 | April 2000 | sediment | <i>Yersinia</i> sp. |
| Canal 1 | April 2000 | water | <i>Yersinia enterocolitica</i> |
| Canal 4 | April 2000 | water | <i>Yersinia intermedia</i> |
| Canal 1 | April 2000 | water | <i>Yersinia intermedia</i> |
| CW 2 | April 2000 | sediment | <i>Yersinia intermedia</i> |
| CW 1 | April 2000 | water | <i>Yersinia enterocolitica</i> |
| CW 1 | April 2000 | water | <i>Yersinia enterocolitica</i> |
| CW 2 | April 2000 | water | <i>Yersinia intermedia</i> |
| CW 2 | April 2000 | water | <i>Yersinia intermedia</i> |
| CW 2 | April 2000 | water | <i>Yersinia</i> sp / <i>Aeromonas</i> sp. |
| MM 4 | June 2000 | water | <i>Yersinia</i> sp. |
| MM 1 | April 2000 | sediment | <i>Yersinia intermedia</i> |
| MM 1 | April 2000 | sediment | <i>Yersinia intermedia</i> |
| MM 2 | April 2000 | sediment | <i>Yersinia intermedia</i> |
| MM 3 | April 2000 | sediment | <i>Yersinia intermedia</i> |
| MM 3 | April 2000 | sediment | <i>Yersinia</i> sp. |
| MM 1 | April 2000 | water | <i>Yersinia intermedia</i> |
| MM 1 | April 2000 | water | <i>Yersinia intermedia</i> |
| MM 3 | April 2000 | water | <i>Yersinia intermedia</i> |

Note: CW = Hastings Pork created wetland, MM = MM Marsh.

Table A.5. Concentrations (mg/L) of trace elements in water samples collected from Waterfowl Production Areas (federally managed wetlands) in the Rainwater Basin, Clay County, Nebraska, 1992-2001.

| Catalog No. | General Location | Sample ID | Date Collected | | Trace Element Concentration mg/L wet weight | | | | | | |
|-------------|----------------------|-----------|----------------|------|---|-----------|-----------|-----------|-----------|-----------|---------|
| | | | Month | Year | Al | As | B | Ba | Be | Cd | Cr |
| 6050045 | Harvard WPA | M1M93 | August | 1993 | 8.04 | <0.0056 | 0.0361 | 0.0776 | <0.0005 | <0.0005 | <0.005 |
| 6050045 | McMurtrey NWR | M6M93 | August | 1993 | 1.24 | <0.0056 | 0.0292 | 0.0478 | <0.0005 | <0.0005 | <0.005 |
| 6050045 | Massie WPA | W10M93 | August | 1993 | 0.119 | <0.0056 | 0.0577 | 0.252 | <0.0005 | <0.0005 | <0.005 |
| 6050045 | Harvard WPA | W1M93 | August | 1993 | 0.154 | <0.0056 | 0.005 | 0.15 | <0.0005 | <0.0005 | <0.005 |
| 6050045 | Harvard WPA | W4M93 | August | 1993 | 0.168 | <0.0056 | 0.0458 | 0.124 | <0.0005 | <0.0005 | <0.005 |
| 6050045 | McMurtrey NWR | W5M93 | August | 1993 | 0.183 | <0.0056 | 0.0718 | 0.19 | <0.0005 | <0.0005 | <0.005 |
| 6050045 | McMurtrey NWR | W6M93 | August | 1993 | 0.119 | <0.0055 | 0.0412 | 0.223 | 0.0023 | 0.0019 | 0.0089 |
| 6050045 | McMurtrey NWR | W7M93 | August | 1993 | 0.091 | <0.0056 | 0.0375 | 0.218 | 0.0012 | 0.0009 | 0.005 |
| 6050045 | Massie WPA | W9M93 | August | 1993 | 0.509 | 0.0088 | 0.0536 | 0.192 | 0.0032 | 0.0036 | 0.0075 |
| 6050017 | Harvard or McMurtrey | M1M | October | 1991 | NC | 0.00202 | NC | NC | NC | NC | NC |
| 6050017 | Harvard or McMurtrey | M6M | October | 1991 | NC | 0.00258 | NC | NC | NC | NC | NC |
| 6050017 | Harvard or McMurtrey | W1M | October | 1991 | NC | 0.000619 | NC | NC | NC | NC | NC |
| 6050017 | Harvard or McMurtrey | W4M | October | 1991 | NC | 0.00146 | NC | NC | NC | NC | NC |
| 6050017 | Harvard or McMurtrey | W5M | October | 1991 | NC | 0.00126 | NC | NC | NC | NC | NC |
| 6050017 | Harvard or McMurtrey | W6M | October | 1991 | NC | 0.00154 | NC | NC | NC | NC | NC |
| 6050017 | Harvard or McMurtrey | W7M | October | 1991 | NC | 0.00134 | NC | NC | NC | NC | NC |
| 6050062 | McMurtrey NWR | 320ME1 | March | 2000 | 0.657 | <0.0056 | 0.047 | 0.0827 | <0.0011 | <0.0011 | <0.0056 |
| 6050066 | McMurtrey NWR | MW | October | 2000 | 11.9 | 0.0119 | <0.111 | 0.59 | 0.0019 | 0.0014 | 0.0113 |
| | | | | | Cu | Fe | Hg | Mg | Mn | Mo | |
| 6050045 | Harvard WPA | M1M93 | August | 1993 | <0.005 | <0.02 | <0.002 | 2.46 | <0.002 | <0.004 | |
| 6050045 | McMurtrey NWR | M6M93 | August | 1993 | <0.005 | 5.81 | <0.002 | 2.01 | <0.002 | <0.004 | |
| 6050045 | Massie WPA | W10M93 | August | 1993 | <0.005 | 1.18 | <0.002 | 7.85 | 0.0817 | <0.004 | |
| 6050045 | Harvard WPA | W1M93 | August | 1993 | <0.005 | 0.301 | <0.002 | 12.2 | 0.085 | <0.004 | |
| 6050045 | Harvard WPA | W4M93 | August | 1993 | <0.005 | 0.0654 | <0.002 | 8.69 | 0.002 | <0.004 | |
| 6050045 | McMurtrey NWR | W5M93 | August | 1993 | <0.005 | 0.345 | <0.002 | 18.9 | 0.0048 | <0.004 | |
| 6050045 | McMurtrey NWR | W6M93 | August | 1993 | 0.0064 | 0.0457 | <0.002 | 8.42 | 0.0024 | 0.0094 | |
| 6050045 | McMurtrey NWR | W7M93 | August | 1993 | 0.0572 | 0.398 | <0.002 | 6.53 | 0.0076 | 0.0062 | |
| 6050045 | Massie WPA | W9M93 | August | 1993 | 0.008 | 0.66 | <0.002 | 11.2 | 0.0457 | 0.0079 | |
| 6050017 | Harvard or McMurtrey | M1M | October | 1991 | NC | NC | <0.00498 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | M6M | October | 1991 | NC | NC | <0.00497 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W1M | October | 1991 | NC | NC | <0.00497 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W4M | October | 1991 | NC | NC | <0.00498 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W5M | October | 1991 | NC | NC | <0.00497 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W6M | October | 1991 | NC | NC | <0.00498 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W7M | October | 1991 | NC | NC | <0.00499 | NC | NC | NC | |
| 6050062 | McMurtrey NWR | 320ME1 | March | 2000 | 0.0062 | 0.398 | <0.0002 | 9.46 | 0.01 | <0.0056 | |
| 6050066 | McMurtrey NWR | MW | October | 2000 | 0.0302 | 12.2 | <0.0002 | 11.5 | 0.85 | <0.0556 | |
| | | | | | Ni | Pb | Se | Sr | V | Zn | |
| 6050045 | Harvard WPA | M1M93 | August | 1993 | <0.005 | <0.005 | <0.0056 | 0.047 | <0.004 | <0.01 | |
| 6050045 | McMurtrey NWR | M6M93 | August | 1993 | <0.005 | <0.005 | <0.0056 | 0.042 | <0.004 | <0.01 | |
| 6050045 | Massie WPA | W10M93 | August | 1993 | <0.005 | <0.005 | 0.0057 | 0.242 | 0.0138 | <0.01 | |
| 6050045 | Harvard WPA | W1M93 | August | 1993 | <0.005 | <0.005 | 0.0071 | 0.429 | 0.0092 | <0.01 | |
| 6050045 | Harvard WPA | W4M93 | August | 1993 | <0.005 | <0.005 | 0.0073 | 0.282 | 0.0054 | <0.01 | |
| 6050045 | McMurtrey NWR | W5M93 | August | 1993 | <0.005 | 0.0056 | 0.0199 | 0.542 | 0.0195 | <0.01 | |
| 6050045 | McMurtrey NWR | W6M93 | August | 1993 | 0.0154 | 0.0053 | 0.0125 | 0.286 | 0.0081 | <0.01 | |
| 6050045 | McMurtrey NWR | W7M93 | August | 1993 | 0.0056 | 0.0101 | 0.0075 | 0.221 | 0.0064 | 0.0274 | |
| 6050045 | Massie WPA | W9M93 | August | 1993 | 0.0172 | 0.0175 | 0.0062 | 0.322 | 0.0183 | 0.0258 | |
| 6050017 | Harvard or McMurtrey | M1M | October | 1991 | NC | NC | 0.00222 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | M6M | October | 1991 | NC | NC | 0.00166 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W1M | October | 1991 | NC | NC | 0.00212 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W4M | October | 1991 | NC | NC | 0.00282 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W5M | October | 1991 | NC | NC | 0.0057 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W6M | October | 1991 | NC | NC | 0.00218 | NC | NC | NC | |
| 6050017 | Harvard or McMurtrey | W7M | October | 1991 | NC | NC | 0.0013 | NC | NC | NC | |
| 6050062 | McMurtrey NWR | 320ME1 | March | 2000 | <0.0056 | <0.0056 | <0.0056 | 0.19 | <0.0056 | <0.0111 | |
| 6050066 | McMurtrey NWR | MW | October | 2000 | 0.017 | 0.0448 | <0.0044 | 0.226 | 0.026 | 0.0881 | |

Note: < indicates the sample was below the detection limit (value = detection limit) and NC = not collected.

Table A.6. Concentrations (mg/kg) of trace elements in sediment samples collected from Waterfowl Production Areas (federally managed wetlands) in the Rainwater Basin, Clay County, Nebraska, 1992-2001.

| Catalog ID | General Location | Sample ID | Date Collected | | Trace Element Concentration in mg/kg dry weight | | | | | | |
|------------|------------------|-----------|----------------|------|---|-------|---------|-------|-------|---------|-------|
| | | | Month | Year | Al | As | B | Ba | Be | Cd | Cr |
| 6050045 | Harvard WPA | HVSM193 | August | 1993 | 10970 | 3.19 | < 4.97 | 175 | 0.73 | < 0.198 | 12.90 |
| 6050045 | Harvard WPA | HVSM293 | August | 1993 | 14360 | 4.57 | < 4.95 | 277 | 0.84 | < 0.198 | 19.20 |
| 6050045 | Harvard WPA | HVSM493 | August | 1993 | 12310 | 3.14 | < 4.96 | 259 | 0.51 | < 0.198 | 14.40 |
| 6050045 | McMurtrey NWR | MCSM593 | August | 1993 | 8386 | 1.81 | < 4.91 | 132 | 0.22 | 0.21 | 12.20 |
| 6050045 | McMurtrey NWR | MCSM693 | August | 1993 | 10330 | 4.35 | 12.20 | 275 | 0.34 | 0.38 | 14.20 |
| 6050045 | McMurtrey NWR | MCSM893 | August | 1993 | 7804 | 2.05 | 7.59 | 114 | 0.92 | 0.62 | 9.56 |
| 6050027 | Eckhart WPA | RWB_E_S | June | 1992 | 11084 | 4.67 | NC | NC | NC | 0.68 | 16.01 |
| 6050027 | Harvard WPA | RWB_H_SM | June | 1992 | 16343 | 4.94 | NC | NC | NC | 0.57 | 26.02 |
| 6050027 | Harvard WPA | RWB_H_SS | June | 1992 | 10811 | 3.26 | NC | NC | NC | 0.25 | 13.15 |
| 6050027 | Smith WPA | RWB_S_S | June | 1992 | 17788 | 4.74 | NC | NC | NC | 0.26 | 14.82 |
| 6050066 | McMurtrey NWR | MS | October | 2000 | 6539 | 2.07 | 3.58 | 227 | 1.00 | 1.32 | 11.70 |
| 6050062 | McMurtrey NWR | 320ME1S | March | 2000 | 12050 | 4.41 | 8.15 | 158 | 0.58 | 0.75 | 11.80 |
| | | | | | Cu | Fe | Hg | Mg | Mn | Mo | |
| 6050045 | Harvard WPA | HVSM193 | August | 1993 | 12.90 | 9850 | < .0994 | 2359 | 400 | < 4.97 | |
| 6050045 | Harvard WPA | HVSM293 | August | 1993 | 16.00 | 13920 | < .0982 | 3481 | 675 | < 4.91 | |
| 6050045 | Harvard WPA | HVSM493 | August | 1993 | 13.60 | 11640 | < .0990 | 3246 | 310 | < 4.95 | |
| 6050045 | McMurtrey NWR | MCSM593 | August | 1993 | 11.70 | 7546 | < .0992 | 2180 | 179 | < 4.96 | |
| 6050045 | McMurtrey NWR | MCSM693 | August | 1993 | 13.90 | 12980 | < .0982 | 3695 | 296 | < 4.91 | |
| 6050045 | McMurtrey NWR | MCSM893 | August | 1993 | 9.54 | 7407 | < .0988 | 1945 | 271 | < 4.94 | |
| 6050027 | Eckhart WPA | RWB_E_S | June | 1992 | 20.33 | 24495 | <0.1 | NC | 345 | NC | |
| 6050027 | Harvard WPA | RWB_H_SM | June | 1992 | 22.82 | 30274 | <0.1 | NC | 316 | NC | |
| 6050027 | Harvard WPA | RWB_H_SS | June | 1992 | 14.41 | 19713 | <0.1 | NC | 240 | NC | |
| 6050027 | Smith WPA | RWB_S_S | June | 1992 | 13.24 | 22971 | <0.1 | NC | 358 | NC | |
| 6050066 | McMurtrey NWR | MS | October | 2000 | 16.20 | 9543 | 0.05 | 2428 | 209 | *5.09 | |
| 6050062 | McMurtrey NWR | 320ME1S | March | 2000 | 14.80 | 10850 | 0.04 | 2278 | 133 | *5.42 | |
| | | | | | Ni | Pb | Se | Sr | V | Zn | |
| 6050045 | Harvard WPA | HVSM193 | August | 1993 | 13.40 | 23.60 | < .497 | 27.70 | 17.30 | 42.50 | |
| 6050045 | Harvard WPA | HVSM293 | August | 1993 | 17.50 | 21.60 | < .491 | 35.50 | 20.70 | 52.20 | |
| 6050045 | Harvard WPA | HVSM493 | August | 1993 | 13.00 | 17.00 | < .495 | 33.90 | 16.40 | 43.00 | |
| 6050045 | McMurtrey NWR | MCSM593 | August | 1993 | 8.54 | 14.20 | < .491 | 31.00 | 12.20 | 36.60 | |
| 6050045 | McMurtrey NWR | MCSM693 | August | 1993 | 14.60 | 13.10 | < .494 | 36.60 | 16.90 | 50.40 | |
| 6050045 | McMurtrey NWR | MCSM893 | August | 1993 | 11.60 | 17.40 | 0.87 | 25.20 | 15.20 | 29.30 | |
| 6050027 | Eckhart WPA | RWB_E_S | June | 1992 | 11.06 | 22.91 | <1 | NC | NC | 101.5 | |
| 6050027 | Harvard WPA | RWB_H_SM | June | 1992 | 22.99 | 30.52 | 1.47 | NC | NC | 125 | |
| 6050027 | Harvard WPA | RWB_H_SS | June | 1992 | 19.48 | 17.45 | <1 | NC | NC | 73.36 | |
| 6050027 | Smith WPA | RWB_S_S | June | 1992 | 18.06 | 15.33 | <1 | NC | NC | 77.27 | |
| 6050066 | McMurtrey NWR | MS | October | 2000 | 11.80 | 22.40 | 1.22 | 47.10 | 25.20 | 63.00 | |
| 6050062 | McMurtrey NWR | 320ME1S | March | 2000 | 10.90 | 20.30 | 1.33 | 32.90 | 19.90 | 57.50 | |

Note: < indicates the sample was below the detection limit (value = detection limit) and NC = not collected.

Table A.7. Antibiotic concentrations in sediment (ng/g) and water (µg/g) from Hastings Pork created wetlands, canals and lagoons, Clay County, Nebraska, 2000.

| Sample ID | Date Col. | Site | Sample Type | Antibiotic concentration (ng/g for sediments and ug/L for water) | | | | | | |
|-----------|-----------|-----------------|-------------|--|-------------------|-----------------|------------|---------|--------------|----------|
| | | | | Tetracycline | Chlortetracycline | Oxytetracycline | Lincomycin | Tylosin | Erythromycin | Tiamulin |
| A127 | 000314 | Lagoon | Water | <10 | <10 | 13 | <2 | <2 | <2 | 4.9 |
| A63 | 000314 | Lagoon | Water | <10 | <10 | 317 | <2 | <2 | <2 | 17.7 |
| C124 | 000314 | Lagoon | Water | 10 | 11 | 318 | <2 | <2 | <2 | 32.6 |
| C416 | 000314 | Lagoon | Water | <10 | 10 | 241 | <2 | <2 | <2 | 13.5 |
| D327 | 000314 | Lagoon | Water | <10 | <10 | 90 | <2 | <2 | <2 | 7.6 |
| D415 | 000314 | Lagoon | Water | <10 | <10 | 282 | <2 | <2 | <2 | 7 |
| D618 | 000314 | Lagoon | Water | <10 | <10 | 77 | <2 | <2 | <2 | 4.9 |
| A127 | 000613 | Lagoon | Water | <10 | <10 | 18.1 | <2 | <2 | <2 | 2.7 |
| A21 | 000613 | Lagoon | Water | 10 | 12 | 426 | 53.4 | <2 | <2 | 5.1 |
| C124 | 000613 | Lagoon | Water | 15 | 18 | 533 | 780.1 | 11.4 | <2 | 3.4 |
| D327 | 000613 | Lagoon | Water | <10 | <10 | 265 | <2 | <2 | <2 | 6.7 |
| A130 | 001003 | Lagoon | Sediment | 186 | 2787 | 7708 | <20 | <20 | <20 | 78 |
| C124 | 001003 | Lagoon | Sediment | 1328 | 6430 | 25549 | <20 | <20 | <20 | 27 |
| D327 | 001003 | Lagoon | Sediment | 119 | 311 | 8943 | <20 | <20 | <20 | <20 |
| A130 | 001003 | Lagoon | Water | <10 | <10 | 13 | <2 | <2 | <2 | 3.4 |
| C124 | 001003 | Lagoon | Water | <10 | 17 | 259 | <2 | <2 | <2 | 6.8 |
| D327 | 001003 | Lagoon | Water | <10 | <10 | 208 | <2 | <2 | <2 | 6.5 |
| A127 | 000314 | Canal | Water | <10 | <10 | 31 | <2 | <2 | <2 | 9.8 |
| C124 | 000314 | Canal | Water | <10 | <10 | 10 | <2 | <2 | <2 | 3.5 |
| D327 | 000314 | Canal | Water | <10 | <10 | 63 | <2 | <2 | <2 | 2.1 |
| A127 | 000613 | Canal | Water | <10 | <10 | 26 | <2 | <2 | <2 | 8.6 |
| C124 | 000613 | Canal | Water | <10 | <10 | 15 | <2 | <2 | <2 | 2.6 |
| D327 | 000613 | Canal | Water | <10 | <10 | <10 | <2 | <2 | <2 | 2.8 |
| A130 | 001003 | Canal | Sediment | 98 | 1800 | 1568 | <20 | <20 | <20 | 29 |
| C124 | 001003 | Canal | Sediment | <30 | 165 | 851 | <20 | <20 | <20 | <20 |
| D327 | 001003 | Canal | Sediment | 50 | 108 | 3494 | <20 | <20 | <20 | <20 |
| A130 | 001003 | Canal | Water | <10 | <10 | 28 | <2 | <2 | <2 | 10.9 |
| C124 | 001003 | Canal | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| D327 | 001003 | Canal | Water | <10 | <10 | <10 | <2 | <2 | <2 | 2.1 |
| D227 | 000613 | Feeder ditch | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| CW1 | 000314 | CW | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| CW5 | 000314 | CW | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| CW1 | 000613 | CW | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| CW5 | 000613 | CW | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| CW1 | 001003 | CW | Sediment | <30 | <30 | 41 | <20 | <20 | <20 | <20 |
| CW5 | 001003 | CW | Sediment | <30 | <30 | <30 | <20 | <20 | <20 | <20 |
| CW1 | 001003 | CW | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| CW5 | 001003 | CW | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| MM | 000314 | McMurtrey Marsh | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |
| MM | 001003 | McMurtrey Marsh | Sediment | <30 | <30 | <30 | <20 | <20 | <20 | <20 |
| MM | 001003 | McMurtrey Marsh | Water | <10 | <10 | <10 | <2 | <2 | <2 | <2 |

Note: Date Col. = the date of sample collection; < indicates below the detection limit (the value is the detection limit) and CW = created wetland.

Table A.8. Concentrations ($\mu\text{g/g}$) of trace elements in invertebrate samples collected from Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000 – 2001.

| Date Col. | Site | Sample ID | Family | Al | | As | | B | | Ba | | Be | | Cd | | Cr | |
|-----------|-------|-----------|--------------|------|------|-------|--------|-------|--------|-------|-------|--------|--------|------|------|-------|------|
| | | | | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. |
| 07/01 | CANAL | C118CI | Chironomidae | 3553 | 443 | 0.935 | 0.117 | 7.14 | 0.89 | 53.30 | 6.65 | 0.18 | 0.02 | 0.16 | 0.02 | 7.87 | 0.98 |
| 07/01 | CANAL | D230CI | Chironomidae | 3578 | 384 | 1.14 | 0.122 | 9.38 | 1.01 | 40.20 | 4.32 | 0.16 | 0.02 | 0.11 | 0.01 | 5.78 | 0.62 |
| 06/00 | CW1 | 601CW1B | Corixidae | 1296 | 407 | 0.298 | 0.0936 | 3.74 | 1.18 | 14.40 | 4.51 | 0.13 | 0.04 | 0.15 | 0.05 | 2.01 | 0.63 |
| 06/00 | CW1 | 601CW1C | Chironomidae | 5283 | 893 | 1.42 | 0.24 | 10.40 | 1.76 | 63.80 | 10.80 | 0.43 | 0.07 | 0.25 | 0.04 | 6.78 | 1.15 |
| 10/00 | CW1 | CW1C | Chironomidae | 6170 | 932 | 1.46 | 0.22 | 11.20 | 1.69 | 73.00 | 11.00 | 0.31 | 0.05 | 0.32 | 0.05 | 5.08 | 0.77 |
| 07/01 | CW2 | CW2I | Chironomidae | 3998 | 497 | 1.1 | 0.137 | 8.17 | 1.02 | 65.30 | 8.11 | 0.20 | 0.02 | 0.51 | 0.06 | 5.29 | 0.66 |
| 07/01 | CW3 | CW3I | Chironomidae | 7040 | 1070 | 1.84 | 0.28 | 9.35 | 1.42 | 66.00 | 10.00 | 0.30 | 0.05 | 0.73 | 0.11 | 8.81 | 1.34 |
| 07/01 | CW4 | CW4I | Chironomidae | 5485 | 615 | 1.88 | 0.211 | 6.51 | 0.73 | 92.20 | 10.30 | 0.26 | 0.03 | 0.63 | 0.07 | 8.00 | 0.90 |
| 06/00 | CW5 | 602CW5B | Corixidae | 2265 | 695 | 0.438 | 0.134 | 4.75 | 1.46 | 35.70 | 11.00 | 0.21 | 0.06 | 0.23 | 0.07 | 3.58 | 1.10 |
| 06/00 | CW5 | 602CW5C | Chironomidae | 7314 | 1485 | 2.32 | 0.471 | 12.00 | 2.44 | 81.70 | 16.60 | 0.51 | 0.10 | 0.47 | 0.09 | 7.92 | 1.61 |
| 10/00 | CW5 | CW5C | Chironomidae | 4619 | 845 | 1.46 | 0.267 | <4.05 | <0.741 | 71.60 | 13.10 | <0.202 | <0.037 | 0.26 | 0.05 | 12.50 | 2.29 |
| 07/01 | CW5 | CW5I | Chironomidae | 3373 | 426 | 1.08 | 0.136 | 3.56 | 0.45 | 41.60 | 5.25 | 0.17 | 0.02 | 0.35 | 0.04 | 4.85 | 0.61 |
| 07/01 | CW6 | CW6I | Chironomidae | 8650 | 1220 | 1.8 | 0.253 | 8.84 | 1.25 | 85.60 | 12.10 | 0.39 | 0.05 | 0.48 | 0.07 | 9.82 | 1.39 |
| 07/01 | CW7 | CW7I | Chironomidae | 2948 | 379 | 1.23 | 0.158 | 3.54 | 0.46 | 42.10 | 5.41 | 0.14 | 0.02 | 1.04 | 0.13 | 3.97 | 0.51 |

| Date Col. | Site | Sample ID | Family | Cu | | Fe | | Hg | | Mg | | Mn | | Mo | |
|-----------|-------|-----------|--------------|-------|------|------|------|---------|----------|------|------|--------|-------|-------|---------|
| | | | | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. |
| 07/01 | CANAL | C118CI | Chironomidae | 25.90 | 3.24 | 2574 | 321 | <0.2 | <0.025 | 2628 | 328 | 121.00 | 15.10 | <2.00 | <0.25 |
| 07/01 | CANAL | D230CI | Chironomidae | 19.50 | 2.10 | 3219 | 345 | <0.2 | <0.0215 | 2090 | 224 | 62.70 | 6.72 | <2.00 | <0.215 |
| 06/00 | CW1 | 601CW1B | Corixidae | 18.70 | 5.89 | 1154 | 362 | <0.0977 | <0.0307 | 1382 | 434 | 30.30 | 9.50 | <0.48 | <0.151 |
| 06/00 | CW1 | 601CW1C | Chironomidae | 19.70 | 3.33 | 5788 | 978 | <0.0995 | <0.0168 | 2862 | 484 | 126.00 | 21.40 | <0.48 | <0.0812 |
| 10/00 | CW1 | CW1C | Chironomidae | 23.00 | 3.47 | 6718 | 1014 | <0.0408 | <0.00616 | 3268 | 493 | 142.00 | 21.50 | <1.02 | <0.154 |
| 07/01 | CW2 | CW2I | Chironomidae | 15.50 | 1.92 | 3026 | 376 | <0.2 | <0.0248 | 2098 | 261 | 143.00 | 17.80 | <2.00 | <0.248 |
| 07/01 | CW3 | CW3I | Chironomidae | 16.60 | 2.52 | 4750 | 722 | <0.2 | <0.0304 | 2998 | 456 | 114.00 | 17.30 | <2.00 | <0.304 |
| 07/01 | CW4 | CW4I | Chironomidae | 12.70 | 1.43 | 4257 | 478 | <0.2 | <0.0224 | 3113 | 349 | 222.00 | 24.90 | <2.00 | <0.224 |
| 06/00 | CW5 | 602CW5B | Corixidae | 22.10 | 6.80 | 2012 | 618 | <0.0964 | <0.0296 | 1708 | 524 | 87.40 | 26.80 | <0.50 | <0.153 |
| 06/00 | CW5 | 602CW5C | Chironomidae | 22.80 | 4.64 | 7021 | 1425 | <0.096 | <0.0195 | 3183 | 646 | 151.00 | 30.80 | <0.48 | <0.0969 |
| 10/00 | CW5 | CW5C | Chironomidae | 15.50 | 2.84 | 5216 | 955 | 0.05 | 0.01 | 2359 | 432 | 178.00 | 32.50 | <1.01 | <0.185 |
| 07/01 | CW5 | CW5I | Chironomidae | 31.90 | 4.03 | 2853 | 360 | <0.2 | <0.0253 | 1897 | 240 | 87.00 | 11.00 | <2.00 | <0.253 |
| 07/01 | CW6 | CW6I | Chironomidae | 18.20 | 2.57 | 5222 | 737 | <0.2 | <0.0282 | 3321 | 469 | 145.00 | 20.50 | <2.00 | <0.282 |
| 07/01 | CW7 | CW7I | Chironomidae | 12.10 | 1.55 | 2503 | 322 | <0.2 | <0.0257 | 1930 | 248 | 72.30 | 9.30 | <2.00 | <0.257 |

| Date Col. | Site | Sample ID | Family | Ni | | Pb | | Se | | Sr | | V | | Zn | |
|-----------|-------|-----------|--------------|-------|------|------|------|------|------|-------|------|-------|------|--------|-------|
| | | | | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. |
| 07/01 | CANAL | C118CI | Chironomidae | 7.20 | 0.90 | 1.86 | 0.23 | 2.96 | 0.37 | 68.70 | 8.58 | 7.58 | 0.95 | 159.00 | 19.90 |
| 07/01 | CANAL | D230CI | Chironomidae | 6.67 | 0.72 | 2.07 | 0.22 | 4.27 | 0.46 | 24.60 | 2.64 | 8.48 | 0.91 | 104.00 | 11.20 |
| 06/00 | CW1 | 601CW1B | Corixidae | 2.39 | 0.75 | 2.05 | 0.64 | 1.08 | 0.34 | 6.62 | 2.08 | 2.40 | 0.75 | 124.00 | 39.00 |
| 06/00 | CW1 | 601CW1C | Chironomidae | 18.00 | 3.05 | 9.61 | 1.62 | 0.96 | 0.16 | 19.90 | 3.37 | 11.90 | 2.01 | 85.00 | 14.40 |
| 10/00 | CW1 | CW1C | Chironomidae | 9.81 | 1.48 | 6.86 | 1.04 | 3.30 | 0.50 | 29.00 | 4.38 | 11.00 | 1.67 | 94.90 | 14.30 |
| 07/01 | CW2 | CW2I | Chironomidae | 5.50 | 0.68 | 4.05 | 0.50 | 3.34 | 0.41 | 33.60 | 4.17 | 8.11 | 1.01 | 79.80 | 9.91 |
| 07/01 | CW3 | CW3I | Chironomidae | 9.11 | 1.39 | 4.89 | 0.74 | 2.74 | 0.42 | 24.70 | 3.75 | 13.00 | 1.98 | 75.30 | 11.40 |
| 07/01 | CW4 | CW4I | Chironomidae | 5.87 | 0.66 | 4.23 | 0.48 | 2.72 | 0.31 | 39.70 | 4.45 | 10.00 | 1.13 | 69.80 | 7.83 |
| 06/00 | CW5 | 602CW5B | Corixidae | 4.50 | 1.38 | 2.28 | 0.70 | 1.08 | 0.33 | 13.40 | 4.13 | 4.28 | 1.32 | 174.00 | 53.40 |
| 06/00 | CW5 | 602CW5C | Chironomidae | 20.30 | 4.12 | 9.05 | 1.84 | 0.57 | 0.12 | 25.90 | 5.25 | 16.20 | 3.30 | 80.80 | 16.40 |
| 10/00 | CW5 | CW5C | Chironomidae | 8.44 | 1.54 | 5.49 | 1.00 | 2.28 | 0.42 | 21.20 | 3.87 | 9.68 | 1.81 | 75.30 | 13.80 |
| 07/01 | CW5 | CW5I | Chironomidae | 17.40 | 2.20 | 4.83 | 0.61 | 4.87 | 0.62 | 17.20 | 2.18 | 5.34 | 0.67 | 94.80 | 12.00 |
| 07/01 | CW6 | CW6I | Chironomidae | 8.72 | 1.23 | 4.82 | 0.68 | 3.78 | 0.53 | 26.60 | 3.75 | 14.60 | 2.06 | 106.00 | 14.90 |
| 07/01 | CW7 | CW7I | Chironomidae | 3.41 | 0.44 | 2.51 | 0.32 | 2.30 | 0.30 | 15.00 | 1.94 | 5.36 | 0.69 | 67.30 | 8.66 |

< Indicates the sample was below the detection limit (value = detection limit). Date Col. = approximate date of sample collection, CW = created wetland, D.W. = dry weigh, and W.W. = wet weight.

Table A.9. Concentrations ($\mu\text{g/g}$) of trace elements in sediment samples collected from MM Marsh and Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000.

| Date Col. | Site | Sample ID | Al | | As | | B | | Ba | | Be | | Cd | | Cr | |
|-----------|-----------|-----------|-------|-------|------|-------|------|-------|------|------|-------|-------|-------|---------|------|------|
| | | | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. |
| 07/01 | LAGOON | A127LSa | 7451 | 1562 | 3.52 | 0.738 | 11 | 2.3 | 232 | 48.7 | 0.987 | 0.207 | 3.21 | 0.674 | 48.6 | 10.2 |
| 07/01 | LAGOON | A127LSb | 8999 | 1745 | 4.75 | 0.92 | 14.5 | 2.8 | 211 | 40.8 | 1.17 | 0.228 | 4.47 | 0.867 | 54.5 | 10.6 |
| 07/01 | LAGOON | A63LSa | 14808 | 6587 | 8.38 | 3.73 | <10 | <4.45 | 206 | 91.8 | 1.03 | 0.457 | 1.64 | 0.728 | 43 | 19.1 |
| 07/01 | LAGOON | A63LSb | 13353 | 6253 | 6.39 | 2.99 | 13.8 | 6.45 | 200 | 93.7 | 0.685 | 0.321 | 0.606 | 0.284 | 56.2 | 26.3 |
| 07/01 | LAGOON | C124LSa | 7146 | 1168 | 3.42 | 0.558 | 11.9 | 1.94 | 179 | 29.2 | 0.912 | 0.149 | 2.88 | 0.47 | 38 | 6.22 |
| 07/01 | LAGOON | C124LSb | 5209 | 686 | 2.97 | 0.392 | 11.7 | 1.55 | 174 | 23 | 0.892 | 0.117 | 3.43 | 0.452 | 40.2 | 5.3 |
| 07/01 | LAGOON | D327LSa | 20763 | 7219 | 5.16 | 1.79 | 14.3 | 4.97 | 171 | 59.5 | 1.38 | 0.478 | 1.11 | 0.385 | 33 | 11.5 |
| 07/01 | LAGOON | D327LSb | 14362 | 3377 | 5.38 | 1.27 | 15.4 | 3.62 | 191 | 45 | 1.43 | 0.336 | 1.99 | 0.468 | 36.1 | 8.49 |
| 07/01 | LAGOON | D415LSa | 7620 | 1282 | 4.49 | 0.756 | 16.9 | 2.85 | 194 | 32.6 | 1.25 | 0.21 | 2.7 | 0.455 | 39.4 | 6.62 |
| 07/01 | LAGOON | D415LSb | 7193 | 1414 | 3.65 | 0.717 | 16.6 | 3.27 | 188 | 36.9 | 1.08 | 0.213 | 2.16 | 0.424 | 37.4 | 7.36 |
| 03/00 | CANAL | 320CC1S | 9710 | 5020 | 2.19 | 1.13 | 4.35 | 2.25 | 216 | 112 | 0.717 | 0.371 | 0.621 | 0.321 | 12.8 | 6.61 |
| 07/01 | CANAL | C118CSa | 14896 | 6827 | 5.07 | 2.32 | 12 | 5.51 | 239 | 110 | 1.05 | 0.481 | 0.842 | 0.386 | 33.6 | 15.4 |
| 07/01 | CANAL | C118CSb | 14736 | 8084 | 4.83 | 2.65 | 12.6 | 6.91 | 347 | 191 | 0.964 | 0.529 | 0.802 | 0.44 | 33.6 | 18.4 |
| 06/00 | CANAL | 607DC2S | 14213 | 5941 | 2.62 | 1.09 | 7.43 | 3.11 | 174 | 72.7 | 0.907 | 0.379 | 0.214 | 0.0895 | 14.3 | 5.99 |
| 10/00 | CANAL | D230CS | 12070 | 4756 | 3.44 | 1.36 | 14.3 | 5.63 | 167 | 65.8 | 0.279 | 0.11 | 0.544 | 0.214 | 12.2 | 4.82 |
| 07/01 | CANAL | D230CSa | 15054 | 7799 | 4.07 | 2.11 | 11.3 | 5.88 | 197 | 102 | 1.11 | 0.578 | 0.392 | 0.203 | 23.5 | 12.2 |
| 07/01 | CANAL | D230CSb | 13834 | 7126 | 4.19 | 2.16 | <10 | <5.15 | 197 | 102 | 0.932 | 0.48 | 0.282 | 0.145 | 19.3 | 9.92 |
| 03/00 | CW1 | 320CW11S | 7396 | 3062 | 2.22 | 0.919 | 5.22 | 2.16 | 191 | 79.2 | 0.786 | 0.325 | 0.617 | 0.255 | 13.9 | 5.75 |
| 06/00 | CW1 | 607CW12S | 16506 | 7791 | 2.5 | 1.18 | 8.83 | 4.17 | 209 | 98.4 | 0.997 | 0.471 | 0.31 | 0.146 | 15.6 | 7.38 |
| 10/00 | CW1 | CW1S | 11550 | 6895 | 5.17 | 3.09 | 16.4 | 9.79 | 126 | 75.2 | 0.401 | 0.239 | 0.464 | 0.277 | 11.6 | 6.94 |
| 07/01 | CW1 | CW1Sa | 21082 | 20207 | 5.67 | 5.43 | 22.9 | 21.9 | 226 | 217 | 1.31 | 1.26 | 0.33 | 0.316 | 26.7 | 25.6 |
| 07/01 | CW1 | CW1Sb | 19492 | 18073 | 5.48 | 5.08 | 19.9 | 18.4 | 268 | 249 | 1.36 | 1.26 | 0.477 | 0.442 | 26.7 | 24.7 |
| 07/01 | CW2 | CW2Sa | 15208 | 5660 | 4.43 | 1.65 | 11.4 | 4.26 | 144 | 53.6 | 1.04 | 0.386 | 0.242 | 0.0901 | 19.1 | 7.13 |
| 07/01 | CW2 | CW2Sb | 20088 | 7555 | 5 | 1.88 | 20.4 | 7.66 | 143 | 53.9 | 1.18 | 0.443 | <0.2 | <0.0752 | 23.6 | 8.89 |
| 07/01 | CW3 | CW3Sa | 21021 | 8585 | 5.76 | 2.35 | 20.1 | 8.2 | 279 | 114 | 1.35 | 0.55 | 0.244 | 0.0996 | 26.6 | 10.9 |
| 07/01 | CW3 | CW3Sb | 19651 | 8004 | 5.37 | 2.19 | 18.7 | 7.62 | 293 | 120 | 1.35 | 0.548 | 0.335 | 0.136 | 26.2 | 10.7 |
| 07/01 | CW4 | CW4Sa | 17304 | 6531 | 4.54 | 1.71 | 19.7 | 7.45 | 197 | 74.4 | 1.01 | 0.382 | <0.2 | <0.0755 | 22.4 | 8.47 |
| 07/01 | CW4 | CW4Sb | 19916 | 7552 | 4.98 | 1.89 | 19.1 | 7.25 | 296 | 112 | 1.51 | 0.572 | 0.281 | 0.107 | 27.2 | 10.3 |
| 03/00 | CW5 | 320CW51S | 7920 | 3453 | 2.42 | 1.06 | 3.13 | 1.36 | 207 | 90.2 | 0.851 | 0.371 | 0.808 | 0.352 | 15 | 6.53 |
| 06/00 | CW5 | 607CW52S | 11279 | 5313 | 1.75 | 0.822 | 9.14 | 4.3 | 154 | 72.3 | 0.748 | 0.352 | 0.255 | 0.12 | 11.7 | 5.5 |
| 10/00 | CW5 | CW5S | 8997 | 5173 | 4.05 | 2.33 | 13.6 | 7.82 | 123 | 70.8 | 0.257 | 0.148 | 0.536 | 0.308 | 9.83 | 5.65 |
| 07/01 | CW5 | CW5Sa | 23280 | 8264 | 5.27 | 1.87 | 22.8 | 8.1 | 285 | 101 | 1.55 | 0.551 | 0.366 | 0.13 | 29.7 | 10.5 |
| 07/01 | CW5 | CW5Sb | 22761 | 9280 | 5.17 | 2.11 | 21.2 | 8.65 | 357 | 146 | 1.51 | 0.615 | 0.543 | 0.221 | 29.5 | 12 |
| 07/01 | CW6 | CW6Sa | 21811 | 7859 | 4.42 | 1.59 | 18 | 6.5 | 250 | 90.2 | 1.45 | 0.523 | 0.333 | 0.12 | 27.5 | 9.9 |
| 07/01 | CW6 | CW6Sb | 22373 | 7453 | 5.22 | 1.74 | 18.1 | 6.03 | 263 | 87.5 | 1.49 | 0.497 | 0.277 | 0.0923 | 26.9 | 8.96 |
| 07/01 | CW7 | CW7Sa | 12601 | 7077 | 2.91 | 1.63 | 12.7 | 7.14 | 143 | 80.2 | 0.886 | 0.497 | <0.2 | <0.112 | 17.7 | 9.92 |
| 07/01 | CW7 | CW7Sb | 14467 | 7273 | 3.63 | 1.83 | 17.3 | 8.67 | 139 | 70.1 | 0.887 | 0.446 | <0.2 | <0.101 | 19.2 | 9.64 |
| 03/00 | McMurtrey | 320ME1S | 6539 | 2570 | 2.07 | 0.814 | 3.58 | 1.41 | 227 | 89.1 | 0.995 | 0.391 | 1.32 | 0.517 | 11.7 | 4.61 |
| 10/00 | McMurtrey | MS | 12050 | 6266 | 4.41 | 2.29 | 8.15 | 4.24 | 158 | 81.9 | 0.581 | 0.302 | 0.748 | 0.389 | 11.8 | 6.11 |

Table A.9. Continued.

| Date | Col. | Site | Sample ID | Cu | | Fe | | Hg | | Mg | | Mn | | Mo | |
|-------|-----------|----------|-----------|------|-------|-------|---------|----------|-------|------|------|------|--------|--------|------|
| | | | | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. |
| 07/01 | LAGOON | A127LSa | 372 | 78 | 8437 | 1768 | <0.2 | <0.0419 | 18181 | 3811 | 1241 | 260 | 10.5 | 2.21 | |
| 07/01 | LAGOON | A127LSb | 406 | 78.7 | 10717 | 2078 | <0.2 | <0.0388 | 21473 | 4164 | 926 | 180 | 11.2 | 2.17 | |
| 07/01 | LAGOON | A63LSa | 109 | 48.7 | 27752 | 12344 | <0.2 | <0.089 | 4364 | 1941 | 968 | 431 | <5.0 | <2.22 | |
| 07/01 | LAGOON | A63LSb | 70 | 32.8 | 12085 | 5659 | <0.2 | <0.0937 | 6675 | 3126 | 2105 | 986 | <5.0 | <2.34 | |
| 07/01 | LAGOON | C124LSa | 301 | 49.1 | 9607 | 1570 | <0.2 | <0.0327 | 9799 | 1601 | 933 | 152 | 13.9 | 2.27 | |
| 07/01 | LAGOON | C124LSb | 351 | 46.2 | 8302 | 1093 | <0.2 | <0.0263 | 10992 | 1448 | 922 | 121 | 17.8 | 2.35 | |
| 07/01 | LAGOON | D327LSa | 204 | 70.9 | 18912 | 6576 | <0.2 | <0.0695 | 12200 | 4242 | 503 | 175 | 6.05 | 2.1 | |
| 07/01 | LAGOON | D327LSb | 439 | 103 | 15667 | 3683 | <0.2 | <0.047 | 17803 | 4186 | 728 | 171 | 14.1 | 3.32 | |
| 07/01 | LAGOON | D415LSa | 598 | 101 | 12992 | 2187 | <0.2 | <0.0337 | 23574 | 3967 | 924 | 156 | 22 | 3.7 | |
| 07/01 | LAGOON | D415LSb | 398 | 78.3 | 11393 | 2240 | <0.2 | <0.0393 | 25057 | 4926 | 997 | 196 | 16.8 | 3.3 | |
| 03/00 | CANAL | 320CC1S | 25.1 | 13 | 10340 | 5346 | <0.0198 | <0.0102 | 7139 | 3691 | 294 | 152 | <4.95 | <2.56 | |
| 07/01 | CANAL | C118CSa | 76.7 | 35.2 | 14134 | 6477 | <0.2 | <0.0917 | 8970 | 4111 | 624 | 286 | <5.0 | <2.29 | |
| 07/01 | CANAL | C118CSb | 68.7 | 37.7 | 15297 | 8392 | <0.2 | <0.11 | 8664 | 4753 | 963 | 528 | <5.0 | <2.74 | |
| 06/00 | CANAL | 607DC2S | 18.9 | 7.88 | 14014 | 5858 | <0.0986 | <0.0412 | 4817 | 2013 | 181 | 75.5 | <0.465 | <0.195 | |
| 10/00 | CANAL | D230CS | 44 | 17.3 | 14260 | 5618 | <0.0214 | <0.00843 | 5079 | 2001 | 294 | 116 | <5.34 | <2.1 | |
| 07/01 | CANAL | D230CSa | 23.9 | 12.4 | 16784 | 8696 | <0.2 | <0.104 | 7841 | 4063 | 394 | 204 | <5.0 | <2.59 | |
| 07/01 | CANAL | D230CSb | 17.7 | 9.1 | 15206 | 7833 | <0.2 | <0.103 | 4585 | 2362 | 304 | 157 | <5.0 | <2.58 | |
| 03/00 | CW1 | 320CW11S | 15.6 | 6.47 | 12040 | 4985 | <0.0202 | <0.00836 | 4333 | 1794 | 354 | 146 | <5.06 | <2.1 | |
| 06/00 | CW1 | 607CW12S | 16.8 | 7.91 | 16148 | 7622 | <0.0676 | <0.0319 | 4583 | 2163 | 322 | 152 | <0.5 | <0.236 | |
| 10/00 | CW1 | CW1S | 13 | 7.76 | 13360 | 7976 | <0.021 | <0.0125 | 3571 | 2132 | 262 | 156 | <5.25 | <3.14 | |
| 07/01 | CW1 | CW1Sa | 25.2 | 24.2 | 20442 | 19594 | <0.2 | <0.192 | 7470 | 7160 | 403 | 387 | <5.0 | <4.79 | |
| 07/01 | CW1 | CW1Sb | 26.2 | 24.3 | 19310 | 17904 | <0.2 | <0.185 | 7280 | 6750 | 523 | 485 | <5.0 | <4.64 | |
| 07/01 | CW2 | CW2Sa | 12 | 4.47 | 16727 | 6226 | <0.2 | <0.0744 | 5293 | 1970 | 231 | 86.1 | <5.0 | <1.86 | |
| 07/01 | CW2 | CW2Sb | 13.3 | 4.99 | 19302 | 7259 | <0.2 | <0.0752 | 6392 | 2404 | 220 | 82.8 | <5.0 | <1.88 | |
| 07/01 | CW3 | CW3Sa | 20.2 | 8.24 | 19987 | 8163 | <0.2 | <0.0817 | 6895 | 2816 | 638 | 261 | <5.0 | <2.04 | |
| 07/01 | CW3 | CW3Sb | 20.8 | 8.49 | 18094 | 7369 | <0.2 | <0.0815 | 7066 | 2878 | 716 | 291 | <5.0 | <2.04 | |
| 07/01 | CW4 | CW4Sa | 12.8 | 4.83 | 17410 | 6571 | <0.2 | <0.0755 | 6198 | 2339 | 456 | 172 | <5.0 | <1.89 | |
| 07/01 | CW4 | CW4Sb | 19.5 | 7.39 | 21986 | 8337 | <0.2 | <0.0758 | 6699 | 2540 | 791 | 300 | <5.0 | <1.9 | |
| 03/00 | CW5 | 320CW51S | 14.8 | 6.43 | 12270 | 5350 | 0.0249 | 0.0109 | 4271 | 1862 | 481 | 210 | <5.04 | <2.2 | |
| 06/00 | CW5 | 607CW52S | 10.6 | 5 | 10841 | 5106 | <0.0619 | <0.0291 | 3475 | 1637 | 302 | 142 | <0.455 | <0.214 | |
| 10/00 | CW5 | CW5S | 10.4 | 5.99 | 10420 | 5992 | <0.021 | <0.0121 | 3566 | 2050 | 230 | 132 | <5.25 | <3.02 | |
| 07/01 | CW5 | CW5Sa | 20 | 7.11 | 21806 | 7741 | <0.2 | <0.071 | 6889 | 2445 | 643 | 228 | <5.0 | <1.78 | |
| 07/01 | CW5 | CW5Sb | 20.1 | 8.18 | 21314 | 8690 | <0.2 | <0.0815 | 6659 | 2715 | 1035 | 422 | <5.0 | <2.04 | |
| 07/01 | CW6 | CW6Sa | 17.8 | 6.41 | 21190 | 7635 | <0.2 | <0.0721 | 7404 | 2668 | 404 | 146 | <5.0 | <1.8 | |
| 07/01 | CW6 | CW6Sb | 19.2 | 6.4 | 21742 | 7242 | <0.2 | <0.0666 | 7379 | 2458 | 394 | 131 | <5.0 | <1.67 | |
| 07/01 | CW7 | CW7Sa | 8.76 | 4.92 | 13353 | 7499 | <0.2 | <0.112 | 4728 | 2655 | 307 | 172 | <5.0 | <2.81 | |
| 07/01 | CW7 | CW7Sb | 9.41 | 4.73 | 14287 | 7182 | <0.2 | <0.101 | 5298 | 2663 | 291 | 146 | <5.0 | <2.51 | |
| 03/00 | McMurtrey | 320ME1S | 16.2 | 6.38 | 9543 | 3750 | 0.0475 | 0.0187 | 2428 | 954 | 209 | 82.1 | <5.09 | <2.0 | |
| 10/00 | McMurtrey | MS | 14.8 | 7.7 | 10850 | 5642 | 0.0406 | 0.0211 | 2278 | 1185 | 133 | 69.4 | <5.42 | <2.82 | |

Table A.9. Continued.

| Date Col. | Site | Sample ID | Ni | | Pb | | Se | | Sr | | V | | Zn | |
|-----------|-----------|-----------|------|------|------|--------|--------|--------|------|------|------|------|------|------|
| | | | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. | D.W. | W.W. |
| 07/01 | LAGOON | A127LSa | 30.7 | 6.43 | 7.19 | 1.51 | 6.78 | 1.42 | 396 | 82.9 | 28.8 | 6.04 | 1712 | 359 |
| 07/01 | LAGOON | A127LSb | 32.9 | 6.39 | 8.77 | 1.7 | 7.86 | 1.52 | 240 | 46.6 | 32.5 | 6.29 | 2059 | 399 |
| 07/01 | LAGOON | A63LSa | 22.9 | 10.2 | 11.7 | 5.22 | 2.58 | 1.15 | 226 | 100 | 29.7 | 13.2 | 614 | 273 |
| 07/01 | LAGOON | A63LSb | 15.3 | 7.18 | 7.51 | 3.52 | 1.51 | 0.706 | 499 | 233 | 33.9 | 15.9 | 394 | 184 |
| 07/01 | LAGOON | C124LSa | 26.3 | 4.29 | 5.58 | 0.911 | 6.81 | 1.11 | 206 | 33.6 | 29.9 | 4.89 | 1969 | 322 |
| 07/01 | LAGOON | C124LSb | 27.2 | 3.58 | <5.0 | <0.658 | 7.66 | 1.01 | 208 | 27.3 | 28.3 | 3.73 | 2364 | 311 |
| 07/01 | LAGOON | D327LSa | 23.9 | 8.32 | 16.6 | 5.78 | 3.81 | 1.32 | 81.3 | 28.3 | 40.6 | 14.1 | 1390 | 483 |
| 07/01 | LAGOON | D327LSb | 28.9 | 6.8 | 9.72 | 2.29 | 7.06 | 1.66 | 132 | 31.1 | 40.1 | 9.42 | 3101 | 729 |
| 07/01 | LAGOON | D415LSa | 32.3 | 5.44 | 6.37 | 1.07 | 10 | 1.69 | 174 | 29.3 | 37.2 | 6.26 | 4820 | 811 |
| 07/01 | LAGOON | D415LSb | 27.2 | 5.35 | 5.52 | 1.08 | 7.08 | 1.39 | 196 | 38.5 | 33.6 | 6.61 | 2916 | 573 |
| 03/00 | CANAL | 320CC1S | 17.9 | 9.23 | 13.8 | 7.16 | <0.495 | <0.256 | 87.2 | 45.1 | 22.1 | 11.4 | 110 | 56.6 |
| 07/01 | CANAL | C118CSa | 22.9 | 10.5 | 11.8 | 5.4 | 2.76 | 1.26 | 290 | 133 | 32.4 | 14.8 | 386 | 177 |
| 07/01 | CANAL | C118CSb | 23.2 | 12.7 | 11.4 | 6.26 | 2.96 | 1.62 | 430 | 236 | 36.4 | 20 | 318 | 175 |
| 06/00 | CANAL | 607DC2S | 15 | 6.25 | 12.8 | 5.33 | 0.428 | 0.179 | 38.5 | 16.1 | 21.9 | 9.13 | 91.3 | 38.2 |
| 10/00 | CANAL | D230CS | 17.7 | 6.99 | 10.6 | 4.18 | 1.48 | 0.583 | 103 | 40.6 | 20.9 | 8.25 | 206 | 81.3 |
| 07/01 | CANAL | D230CSa | 18.1 | 9.38 | 15.9 | 8.23 | 1.23 | 0.637 | 110 | 56.9 | 25.6 | 13.2 | 120 | 61.9 |
| 07/01 | CANAL | D230CSb | 15.7 | 8.07 | 11.6 | 5.97 | * | <0.515 | 46.6 | 24 | 25.5 | 13.1 | 91.4 | 47.1 |
| 03/00 | CW1 | 320CW11S | 16.2 | 6.69 | 12.6 | 5.2 | <0.506 | <0.21 | 45.3 | 18.7 | 22.9 | 9.46 | 59.6 | 24.7 |
| 06/00 | CW1 | 607CW12S | 17 | 8.03 | 14.9 | 7.03 | 0.31 | 0.146 | 42.6 | 20.1 | 26 | 12.3 | 64.4 | 30.4 |
| 10/00 | CW1 | CW1S | 14.3 | 8.54 | 16.1 | 9.6 | <0.42 | <0.251 | 30.3 | 18.1 | 20.6 | 12.3 | 49.2 | 29.4 |
| 07/01 | CW1 | CW1Sa | 20.6 | 19.7 | 15.4 | 14.8 | <1.0 | <0.958 | 97.7 | 93.6 | 37.5 | 35.9 | 119 | 114 |
| 07/01 | CW1 | CW1Sb | 21.2 | 19.7 | 15.6 | 14.5 | 1.06 | 0.985 | 131 | 121 | 36.2 | 33.6 | 125 | 115 |
| 07/01 | CW2 | CW2Sa | 14.9 | 5.54 | 13 | 4.83 | <1.0 | <0.372 | 42.2 | 15.7 | 28.1 | 10.5 | 56.6 | 21.1 |
| 07/01 | CW2 | CW2Sb | 15.7 | 5.92 | 14.4 | 5.4 | <1.0 | <0.376 | 41.6 | 15.7 | 39.6 | 14.9 | 58 | 21.8 |
| 07/01 | CW3 | CW3Sa | 20.3 | 8.28 | 15.9 | 6.5 | <1.0 | <0.408 | 99.6 | 40.7 | 44.3 | 18.1 | 86.7 | 35 |
| 07/01 | CW3 | CW3Sb | 21.7 | 8.85 | 15.1 | 6.16 | <1.0 | <0.407 | 115 | 46.8 | 41.2 | 16.8 | 94.1 | 38.3 |
| 07/01 | CW4 | CW4Sa | 14.5 | 5.47 | 11.4 | 4.32 | <1.0 | <0.377 | 57.7 | 21.8 | 36.9 | 13.9 | 56.6 | 21.4 |
| 07/01 | CW4 | CW4Sb | 23.2 | 8.81 | 16.4 | 6.22 | <1.0 | <0.379 | 73.7 | 27.9 | 45.3 | 17.2 | 86.2 | 32.3 |
| 03/00 | CW5 | 320CW51S | 16.4 | 7.17 | 17.1 | 7.47 | <0.504 | <0.22 | 47.3 | 20.6 | 24.1 | 10.5 | 58.2 | 25.4 |
| 06/00 | CW5 | 607CW52S | 11.5 | 5.42 | 12.3 | 5.78 | 0.255 | 0.12 | 37 | 17.4 | 18 | 8.45 | 42.8 | 20.2 |
| 10/00 | CW5 | CW5S | 11.5 | 6.6 | 13 | 7.46 | 0.57 | 0.328 | 31.6 | 18.1 | 17.4 | 10 | 36.6 | 21.1 |
| 07/01 | CW5 | CW5Sa | 20.7 | 7.33 | 18.8 | 6.67 | <1.0 | <0.355 | 81.8 | 29 | 44.2 | 15.7 | 94 | 33.4 |
| 07/01 | CW5 | CW5Sb | 20.8 | 8.49 | 18.6 | 7.6 | <1.0 | <0.408 | 112 | 45.6 | 43.7 | 17.8 | 93.3 | 38 |
| 07/01 | CW6 | CW6Sa | 20.1 | 7.25 | 15.9 | 5.72 | <1.0 | <0.36 | 66.1 | 23.8 | 41.8 | 15.1 | 86.3 | 31.1 |
| 07/01 | CW6 | CW6Sb | 21 | 7 | 17 | 5.65 | <1.0 | <0.333 | 64.1 | 21.4 | 42.1 | 14 | 91.4 | 30.4 |
| 07/01 | CW7 | CW7Sa | 12.7 | 7.12 | 12 | 6.71 | <1.0 | <0.562 | 37.6 | 21.1 | 27.7 | 15.6 | 42.3 | 23.8 |
| 07/01 | CW7 | CW7Sb | 12.4 | 6.25 | 10.6 | 5.34 | <1.0 | <0.503 | 45 | 22.6 | 30.8 | 15.5 | 42.6 | 21.4 |
| 03/00 | McMurtrey | 320ME1S | 11.8 | 4.62 | 22.4 | 8.82 | 1.22 | 0.479 | 47.1 | 18.5 | 25.2 | 9.92 | 63 | 24.8 |
| 10/00 | McMurtrey | MS | 10.9 | 5.66 | 20.3 | 10.5 | 1.33 | 0.692 | 32.9 | 17.1 | 19.9 | 10.4 | 57.5 | 29.9 |

< Indicates the sample was below the detection limit (value = detection limit). Date Col. = approximate date of sample collection, CW = created wetland, D.W. = dry weigh, W.W. = wet weight.

Table A.10. Concentrations (mg/L) of trace elements in water samples collected from MM Marsh and Hastings Pork created wetlands and canals, Clay County, Nebraska, 2000.

| Date Col. | Site | Sample ID | Al | As | B | Ba | Be | Cd | Cr | Cu | Fe | Hg |
|-----------|-----------|-----------|--------|---------|--------|--------|---------|---------|---------|--------|-------|---------|
| 03/00 | CANAL | 320CC1 | 0.321 | <0.0056 | 0.373 | 0.0514 | <0.0011 | 0.0013 | <0.0056 | 0.106 | 1.18 | >0.0002 |
| 06/00 | CANAL | 607DC2 | 0.879 | 0.00499 | 0.549 | 0.033 | <0.0003 | <0.0004 | 0.00266 | 0.0227 | 0.796 | >0.0002 |
| 10/00 | CANAL | D230CW | 0.154 | 0.0077 | 0.407 | 0.0326 | <0.0006 | <0.0006 | <0.0056 | 0.0334 | 0.261 | >0.0002 |
| 03/00 | CW1 | 320CW11 | 3.61 | 0.0082 | 0.248 | 0.122 | <0.0011 | <0.0011 | <0.0056 | 0.02 | 3.36 | >0.0002 |
| 06/00 | CW1 | 607CW12 | 2.64 | 0.017 | 0.548 | 0.109 | <0.0003 | <0.0004 | 0.0034 | 0.016 | 1.91 | >0.0002 |
| 10/00 | CW1 | CW1W | 0.173 | 0.0101 | 0.427 | 0.0438 | <0.0006 | <0.0006 | <0.0056 | 0.0205 | 0.198 | >0.0002 |
| 03/00 | CW5 | 320CW51 | 14.7 | 0.0166 | 0.204 | 0.335 | <0.0011 | 0.0014 | 0.0141 | 0.0293 | 13.7 | >0.0002 |
| 06/00 | CW5 | 607CW52 | 8.71 | 0.029 | 0.707 | 0.278 | 0.00069 | <0.0004 | 0.00827 | 0.0277 | 5.15 | >0.0002 |
| 10/00 | CW5 | CW5W | <0.111 | 0.0192 | 0.375 | 0.0726 | <0.0006 | <0.0006 | <0.0056 | 0.0172 | 0.171 | >0.0002 |
| 03/00 | McMurtrey | 320ME1 | 0.657 | <0.0056 | 0.047 | 0.0827 | <0.0011 | <0.0011 | <0.0056 | 0.0062 | 0.398 | >0.0002 |
| 10/00 | McMurtrey | MW | 11.9 | 0.0119 | <0.111 | 0.59 | 0.0019 | 0.0014 | 0.0113 | 0.0302 | 12.2 | >0.0002 |

| Date Col. | Site | Sample ID | Mg | Mn | Mo | Ni | Pb | Se | Sr | V | Zn |
|-----------|-----------|-----------|------|--------|---------|---------|---------|---------|-------|---------|---------|
| 03/00 | CANAL | 320CC1 | 31.6 | 0.307 | 0.0167 | 0.034 | 0.0086 | <0.0056 | 0.442 | 0.006 | 0.566 |
| 06/00 | CANAL | 607DC2 | 51.4 | 0.11 | 0.00533 | 0.0437 | 0.00102 | 0.0035 | 0.261 | 0.0166 | 0.101 |
| 10/00 | CANAL | D230CW | 50.8 | 0.0951 | <0.0556 | 0.0358 | <0.0111 | <0.0044 | 0.22 | 0.006 | 0.103 |
| 03/00 | CW1 | 320CW11 | 25.2 | 0.287 | 0.0073 | 0.0233 | 0.0099 | <0.0056 | 0.244 | 0.0165 | 0.0663 |
| 06/00 | CW1 | 607CW12 | 50.9 | 0.244 | <0.004 | 0.0506 | 0.00295 | 0.0027 | 0.256 | 0.019 | 0.0584 |
| 10/00 | CW1 | CW1W | 52.3 | 0.0856 | <0.0556 | 0.0289 | <0.0111 | <0.0044 | 0.188 | 0.0086 | 0.0399 |
| 03/00 | CW5 | 320CW51 | 31.4 | 0.874 | <0.0556 | 0.0323 | 0.0245 | <0.0056 | 0.261 | 0.0458 | 0.0944 |
| 06/00 | CW5 | 607CW52 | 55.8 | 0.556 | 0.0144 | 0.0739 | 0.00861 | 0.0069 | 0.366 | 0.0399 | 0.0667 |
| 10/00 | CW5 | CW5W | 35 | 0.131 | <0.0556 | 0.0271 | <0.0111 | <0.0044 | 0.182 | 0.0119 | 0.013 |
| 03/00 | McMurtrey | 320ME1 | 9.46 | 0.01 | <0.0556 | <0.0556 | <0.0556 | <0.0556 | 0.19 | <0.0556 | <0.0111 |
| 10/00 | McMurtrey | MW | 11.5 | 0.85 | <0.0556 | 0.017 | 0.0448 | <0.0044 | 0.226 | 0.026 | 0.0881 |

< Indicates the sample was below the detection limit (value = detection limit). Date Col. = approximate date of sample collection, CW = created wetland.